

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
4 September 2003 (04.09.2003)

PCT

(10) International Publication Number
WO 03/073097 A2

- (51) International Patent Classification⁷: G01N 33/50, C07K 14/705, 14/00, A61K 39/00, C12N 5/06
- (74) Agent: SONN & PARTNER; Riemergasse 14, A-1010 Vienna (AT).
- (21) International Application Number: PCT/EP03/02005
- (22) International Filing Date: 27 February 2003 (27.02.2003)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
A 316/2002 28 February 2002 (28.02.2002) AT
A 1376/2002 13 September 2002 (13.09.2002) AT
- (71) Applicant (for all designated States except US): INTERCELL AG [AT/AT]; Campus Vienna Biocenter 6, A-1030 Vienna (AT).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): KLADE, Christoph [AT/AT]; Gröhrmühlgasse 1B/17, A-2700 Wr. Neustadt (AT). SCHALICH, Julianne [AT/NL]; Bolmanstraat 17, NL-2496 RE The Hague (NL). VYTVYTSKA, Oresta [UA/AT]; Leystrasse 110/1/2, A-1200 Vienna (AT). ZAUNER, Wolfgang [AT/AT]; Parkgasse 13/3/22, A-1030 Vienna (AT). BIRSTIEL, Max [CH/CH]; Nucleo, CH-6953 Lugaggia (CH). AICHINGER, Gerald [AT/AT]; Czartoryskigasse 47, A-1180 Vienna (AT). OTAVA, Alexander [UA/AT]; Leystrasse 110/1/2, A-1200 Vienna (AT). MATTNER, Frank [DE/AT]; Krottenbachstrasse 267/D12, A-1190 Vienna (AT).
- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Declarations under Rule 4.17:**
— as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii)) for all designations of inventorship (Rule 4.17(iv)) for US only
- Published:**
— without international search report and to be republished upon receipt of that report
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

WO 03/073097 A2

(54) Title: METHOD FOR ISOLATING LIGANDS

(57) Abstract: Described is a method for isolating ligands which have a binding capacity to a MHC/HLA molecule or a complex comprising said ligand and said MHC/HLA molecule which method comprises the following steps: - providing a pool of ligands, said pool containing ligands which bind to said MHC/HLA molecule and ligands which do not bind to said MHC/HLA molecule, - contacting said MHC/HLA molecule with said pool of ligands whereby a ligand which has a binding capacity to said MHC/HLA molecule binds to said MHC/HLA molecule and a complex comprising said ligand and said MHC/HLA molecule is formed, - detecting and optionally separating said complex from the ligands which do not bind to said MHC/HLA molecule and - optionally isolating and characterising the ligand from said complex as well as a method for isolating T cell epitopes which have a binding capacity to a MHC/HLA molecule.

- 1 -

Method for Isolating Ligands

The present invention relates to a method for isolating ligands, especially for isolating T cell epitopes which have a binding capacity to a MHC/HLA molecule.

The immune system is a complex network of inter-related cell types and molecules, which has evolved in order to protect multicellular organisms from infectious microorganisms. It can be divided into the evolutionary older innate (or natural) immunity and adaptive (or acquired) immunity. The innate immune system recognizes patterns, which are usually common and essential for pathogens. For this limited number of molecular structures germ-line encoded receptors have evolved. By contrast, cells of the adaptive immune system - B and T lymphocytes - can recognize a huge variety of antigenic structures. The receptors, termed according to the cell types expressing them, B cell receptor (BCR, its soluble versions are called antibodies) and T cell receptor (TCR, only cell-surface associated forms) are generated by somatic recombination and show a clonal distribution. Thus, initially there is only small number of cells with a certain specificity. Upon antigen encounter these cells start to divide (clonal expansion) to generate an effector population able to cope with the antigen. After elimination of antigen a specialized sub-population of cells specifically recognizing this antigen remains as immunological memory. Taken together the adaptive immune system is slow (compared to innate immunity), however specific and it improves upon repeated exposure to a given pathogen/antigen.

T cells have a central role in adaptive immunity. Their receptors (TCRs) recognize "major histocompatibility complex" (MHC or HLA):peptide complexes on the surface of cells. These peptides are called T cell epitopes and represent degradation products of antigens. There are two major classes of T cells: CD8-positive cytotoxic T cells (CTL) are restricted to MHC class I. CD4-positive helper T cells (HTL) are restricted to MHC class II. HTL are essential for many features of adaptive immunity: activation of so called "professional antigen-presenting cells" (APCs), immunoglobulin (Ig) class switch, the germinal center reaction and

Ig affinity maturation, activation of CTL, immunological memory, regulation of the immune response and others.

MHC molecules collect peptides inside the cell and present them on the cell surface to TCRs of T cells. There are two major classes of MHC, class I recognized by CD8-positive CTL and class II recognized by CD4-positive HTL.

MHC class I molecules consist of a membrane-anchored alpha-chain of 45 kDa and the non-covalently attached b2-microglobulin (b2m) of 12 kDa. Resolution of the 3-dimensional structure by X-ray crystallography (Stern and Wiley 1994) revealed that the alpha-chain possesses a cleft, which is closed at both ends and accommodates peptides from 8 to 11 amino acids length. Class I molecules are ubiquitously expressed, and the peptides they present originate from cytoplasmic proteins. These are degraded by the proteasome, and the resulting peptides are actively transported into the endoplasmatic reticulum (ER). There, with the help of several chaperones, MHC:peptide complexes are formed and transported to the cell surface (Heemels 1995). Thus, MHC class I mirrors the proteome of a cell on its surface and allows T cells to recognize intracellular pathogens or malignant cells.

MHC class II molecules consist of two membrane-anchored proteins (alpha- and beta-chain) of 35 kDa and 30 kDa, respectively. These together form a cleft, open at both ends, which can accommodate peptides of variable length, usually from 12 to 25 amino acids. Despite these differences, class I and II molecules share surprising structural similarity (Stern and Wiley 1994). Class II molecules are only expressed on professional APC including dendritic cells (DC), B-cells and macrophages/monocytes. These cells are specialized in taking up and processing antigens in the endosomal pathway. Immediately after their biosynthesis, class II molecules are complexed by the so-called invariant chain (Ii), which prevents binding of peptides in the ER. When vesicles containing class II:Ii complexes fuse with endosomes containing degradation products of exogenous antigen, Ii is degraded until the MHC binding cleft is only complexed by the so-called CLIP peptide. The latter is with the help of chaperones like HLA-DM exchanged by antigenic peptides (Villadangos 2000).

Finally, MHC:peptide complexes are again presented on the surface of APCs, which interact in numerous ways with HTL.

Being both polygenic and extremely polymorphic, the MHC system is highly complex. For the class I alpha-chain in humans there are three gene loci termed HLA-A, -B and -C. Likewise, there are three class II alpha-chain loci (DRA, DQA, DPA); for class II beta-chain loci the situation is even more complex as there are four different DR beta-chains (DRB1,2,3,5) plus DQB and DPB. Except the monomorphic DR alpha-chain DRA, each gene locus is present in many different alleles (dozens to hundreds) in the population (Klein 1986). Different alleles have largely distinct binding specificities for peptides. Alleles are designated, for example, HLA-A*0201 or HLA-DRB1*0401 or HLA-DPA*0101/DPB*0401.

T cell epitopes have been identified by a variety of approaches (Van den Eynde 1997). T cell lines and clones have for instance been used to screen cDNA expression libraries for instance in the context of COS cells transfected with the appropriate HLA-molecule. Alternatively, biochemical approaches have been pursued. The latter involved elution of natural ligands from MHC molecules on the surface of target cells, the separation of these peptides by several chromatography steps, analysis of their reactivity with lymphocytes in epitope reconstitution assays and sequencing by mass spectrometry (Wölfel et al. 1994, Cox et al. 1994).

Recently the advent of highly sensitive cytokine detection assays like the IFN- γ ELISpot allowed using lymphocytes directly ex vivo for screening of overlapping synthetic peptides (Maecker 2001, Kern 2000, Tobery 2001). Primarily, Kern et al. (1999&2000) used arrays of pools of overlapping 9mer peptides to map CD8+ T cell epitopes in vitro. Later, Tobery et al., 2001 modified this approach and demonstrated that pools containing as many as 64 20mer peptides may be used to screen for both CD8+ and CD4+ T cell epitopes in mice. Both these methods were based on the monitoring of antigen-specific response by measuring INF- γ production either by intracellular staining (Kern et al 2000) or in ELISpot assay (Tobery et al., 2001). By use of mixtures of 15-mers the CD4+ T cell responses are approximately

- 4 -

equal to those detected when whole soluble protein was used as an antigen, while -not surprising- the CD8+ T cell responses are significantly higher than the often negligible responses detected with soluble protein stimulation. Furthermore, the CD8+ T cell responses to a mixture of 15 amino acid peptides are similar to those obtained with a mix of 8-12 amino acid peptides, selected to represent known MHC class I minimal epitopes. Most probably peptidases associated with the cell membrane are responsible for "clipping" peptides to optimal length under these circumstances (Maecker et al, 2001).

An interesting alternative is to screen synthetic combinatorial peptide libraries with specific lymphocytes. For instance, a decapeptide library consisting of 200 mixtures arranged in a positional scanning format, has been successfully used for identification of peptide ligands that stimulate clonotypic populations of T cells (Wilson, et al., J. Immunol., 1999, 163:6424-6434).

Many T cell epitopes have been identified by so called "Reverse immunological approaches" (Rammensee 1999). In this case the protein giving rise to a potential T cell epitope is known, and its primary sequence is scanned for HLA binding motifs. Typically dozens to hundreds of candidate peptides or even a full set of overlapping peptides are synthesized and tested for binding to HLA molecules. Usually, the best binders are selected for further characterization with regard to their reactivity with T cells. This can for instance be done by priming T cells in vitro or in vivo with the help of HLA transgenic mice.

It is an object of the present invention to provide a method for screening ligands for specific MHC molecules, preferably for delivering suitable and specific T cell epitopes selected from a variety of ligands having unknown specificity for a given MHC molecule.

Therefore the present invention provides a method for isolating ligands which have a binding capacity to a MHC/HLA molecule or a complex comprising said ligand and said MHC/HLA molecule which method comprises the following steps:

- 5 -

- providing a pool of ligands, said pool containing ligands which bind to said MHC/HLA molecule and ligands which do not bind to said MHC/HLA molecule,
- contacting said MHC/HLA molecule with said pool of ligands whereby a ligand which has a binding capacity to said MHC/HLA molecule binds to said MHC/HLA molecule and a complex comprising said ligand and said MHC/HLA molecule is formed,
- detecting and optionally separating said complex from the ligands which do not bind to said MHC/HLA molecule and
- optionally isolating and characterising the ligand from said complex.

The present invention also provides a method for isolating T cell epitopes which have a binding capacity to a MHC/HLA molecule or a complex comprising said epitope and said MHC/HLA molecule which method comprises the following steps:

- providing a pool of ligands, said pool containing ligands which bind to a MHC/HLA molecule and ligands which do not bind to said MHC/HLA molecule,
- contacting said MHC/HLA molecule with said pool of ligands whereby a ligand which has a binding capacity to said MHC/HLA molecule binds to said MHC/HLA molecule and a complex comprising said ligand and said MHC/HLA molecule is formed,
- detecting and optionally separating said complex from the ligands which do not bind to said MHC/HLA molecule,
- optionally isolating and characterising the ligand from said complex,
- assaying said optionally isolated ligand or said complex in a T cell assay for T cell activation capacity and
- providing the optionally isolated ligand with a T cell activation capacity as T cell epitope or as complex.

The method according to the present invention enables a screening system for screening binding capacity to specific MHC/HLA molecules. Identifying MHC binding molecules is an important tool for molecular characterisation of pathogens, tumors, etc. It is therefore possible with the present invention to screen a variety (a "pool") of potential ligands at once for their functional affinity towards MHC molecules. Binding affinity towards MHC molecules is also a necessary prerequisite for ligands in-

- 6 -

tended to be used as T cell epitopes, although not a sufficient one. Suitable T cell epitope candidates have also to be screened and assayed with respect to their T cell activation capacity. The combination of the screening method for binding according to the present invention with a suitable T cell assay therefore provides the method for isolating T cell epitopes according to the present invention wherein such T cell epitopes are identifiable out of a pool of potential ligands using an MHC binding assay.

In contrast to the prior art, where such assays have always been performed on ligands with known binding/MHC specificity, the methods according to the present invention provide such assays as a screening tool for pools with ligands of unknown specificity. In the prior art such assays have been typically performed on individual single ligands, to test their binding affinity to MHC/HLA molecules. In Kwok et al. (2001) pools of maximally up to 5 overlapping synthetic peptides were used to generate MHC class II tetramers; the latter were then used to stain PBMC for T cells specific for particular MHC class II:peptide complexes which were generated in the binding reaction with the pools of 5 peptides. However, an increase in the number of ligands per pool in such an approach was not regarded as being possible, both for sensitivity and specificity reasons (Novak et al. 2001). A problem with regard to specificity would be the generation of MHC tetramers with more than one binder per tetramer, if more than one binder would be present in the pool. This would preclude staining of T cells, which is used for identification of epitopes in the approach described in the prior art. In strong contrast to that the approach according to the present invention allows the identification of more than one binder out of highly complex mixtures containing more than one binder.

The nature of the pool to be screened with the present invention is not critical: the pools may contain any naturally or not naturally occurring substance which a) binds specifically to MHC/HLA molecules and/or b) may be specifically recognized by T cells. The binding properties of the set of ligands of the pool with respect to MHC molecules is not known; therefore, usually

- 7 -

binders and at least a non-binder for a given MHC molecule are contained in the pool. The pool therefore comprises at least ten different ligands. Practically, pools are used according to the present invention containing significantly more different ligand species, e.g. 20 or more, 100 or more, 1.000 or more or 10.000 or more. It is also possible to screen larger libraries (with e.g. more than 10^6 , more than 10^8 or even more than 10^{10} different ligand species). This, however, is mainly dependent on the availability of such ligand libraries.

Evidently, MHC:peptide complexes are not typical receptor-ligand systems and have hitherto not been regarded as being the basis of a proper screening tool. In vivo, usually only MHC:peptide complexes but not empty MHC molecules exist. MHC:peptide complexes are not generated by a simple binding of a peptide to an empty MHC molecule, but through the highly complex - and still not fully understood - process of so called "antigen processing and presentation". This is a highly organized intracellular process involving multiple enzymes (cytosolic and lysosomal proteases, transporters, chaperones, peptide-exchange factors, etc.). In fact it is well known that MHC molecules without ligand are unstable and undergo rapid degradation.

Therefore, a main feature of the present invention is the development of production, purification and reaction conditions providing recombinant "empty" MHC-molecules and enabling their use to "capture" ligands" from pools. There are no examples in the prior art demonstrating a similar possibility. The above cited Novak et al. reference discloses a production (insect cells) and purification strategy to obtain recombinant MHC molecules, which are subsequently incubated with a few peptides and tetramerized. The MHC tetramers are used to stain cells from individuals who are likely to have T-cells against the antigen represented by the peptides, thus providing the necessary proof, that the ligand represents also a true T-cell epitope. However, the mentioned prior art explicitly states that only up to 5 peptides could be used successfully. This is in strong contrast to the present approach, which may be successfully applied for 10, 21 and even several hundreds or thousands

of peptides per pool.

Preferred pools of ligands to be used in the method according to the present invention are selected from the group consisting of a pool of peptides, especially overlapping peptides, a pool of protein fragments, a pool of glycolipids, a pool of glycosphingolipids, a pool of lipopeptides, a pool of lipids, a pool of glycans, a pool of modified peptides, a pool obtained from antigen-presenting cells, preferably in the form of total lysates or fractions thereof, especially fractions eluted from the surface or the MHC/HLA molecules of these cells, a pool comprised of fragments of cells, especially pathogen cells, tumor cells or tissues, a pool comprised of peptide libraries, pools of (poly)peptides generated from recombinant DNA libraries, especially derived from pathogens or tumor cells, a pool of proteins and/or protein fragments from a specific pathogen or mixtures thereof.

The ligands of the pools may be derived from natural sources (in native and/or derivatised form) but also be produced synthetically (e.g. by chemical synthesis or by recombinant technology). If (poly)peptide ligands are provided in the pools, those peptides are preferably generated by peptide synthesizers or by recombinant technology. According to a preferred embodiment, a pool of (poly)peptides may be generated from recombinant DNA libraries, e.g. derived from pathogens or tumor cells, by in vitro translation (e.g. by ribosome display) or by expression through heterologous hosts like E.coli or others.

Ligands are therefore preferably peptides being fragments of antigens which could serve as T cell epitopes. Such peptides should preferably be longer than 6, especially longer than 8 amino acids and have preferred maximum lengths of 40, 30, 20, 15 or even 11 or 12 amino acids.

Preferred pathogens wherefrom such peptides can be taken are selected from human immune deficiency virus (HIV), hepatitis A and B viruses, hepatitis C virus (HCV), Rous sarcoma virus (RSV), Epstein Barr virus (EBV), Influenza virus, Rotavirus, Staphylococcus aureus, Chlamydia pneumoniae, Chlamydia trachomatis, Mycobacterium tuberculosis, Streptococcus pneumoniae, Bacillus

antracis, *Vibrio cholerae*, *Plasmodium* sp. (*Pl. falciparum*, *Pl. vivax*, etc.), *Aspergillus* sp. or *Candida albicans*. Antigens may also be molecules expressed by cancer cells (tumor antigens). In the same way also tumor antigens (cancer vaccines) or autoimmune antigens may be used for providing suitable (peptide) ligands for the present invention.

The nature of the specific MHC molecules (of course also MHC-like molecules are encompassed by this term) to be selected for the present methods is again not critical. Therefore, these molecules may be selected in principle from any species, especially primates like humans (HLA, see below), chimpanzees, other mammals, e.g. macaques, rabbits, cats, dogs or rodents like mice, rats, guinea pigs and others, agriculturally important animals like cattle, horses, sheep and fish, although human (or "humanized") molecules are of course preferred for providing vaccines for humans. For providing vaccines for specific animals, especially agriculturally important animals, like cattle, horses, sheep and fish, the use of MHC molecules being specific for these animals is preferred.

Preferred HLA molecules therefore comprise Class I molecules derived from the HLA-A, -B or -C loci, especially A1, A2, A3, A24, A11, A23, A29, A30, A68; B7, B8, B15, B16, B27, B35, B40, B44, B46, B51, B52, B53; Cw3, Cw4, Cw6, Cw7; Class II molecules derived from the HLA-DP, -DQ or -DR loci, especially DR1, DR2, DR3, DR4, DR7, DR8, DR9, DR11, DR12, DR13, DR51, DR52, DR53; DP2, DP3, DP4; DQ1, DQ3, DQ5, DQ6; and non-classical MHC/HLA and MHC/HLA-like molecules, which can specifically bind ligands, especially HLA-E, HLA-G, MICA, MICB, Qa1, Qa2, T10, T18, T22, M3 and members of the CD1 family.

According to a preferred embodiment, the methods according to the present invention is characterised in that said MHC/HLA molecules are selected from HLA class I molecules, HLA class II molecules, non classical MHC/HLA and MHC/HLA-like molecules or mixtures thereof, or mixtures thereof.

Preferably, the optional characterising step of the ligands of the complex is performed by using a method selected from the

group consisting of mass spectroscopy, polypeptide sequencing, binding assays, especially SDS-stability assays, identification of ligands by determination of their retention factors by chromatography, especially HPLC, or other spectroscopic techniques, especially violet (UV), infra-red (IR), nuclear magnetic resonance (NMR), circular dichroism (CD) or electron spin resonance (ESR), or combinations thereof.

According to a preferred embodiment the method of the present invention is characterised in that it is combined with a cytokine secretion assay, preferably with an Elispot assay, an intracellular cytokine staining, FACS or an ELISA (enzyme-linked immunoassays) (see e.g. Current Protocols in Immunology).

Preferred T cell assays comprise the mixing and incubation of said complex with isolated T cells and subsequent measuring cytokine secretion or proliferation of said isolated T cells and/or the measuring up-regulation of activation markers, especially CD69, CD38, or down-regulation of surface markers, especially CD3, CD8 or TCR and/or the measuring up-/down-regulation of mRNAs involved in T cell activation, especially by real-time RT-PCR (see e.g. Current Protocols in Immunology, Current Protocols in Molecular Biology). The T cell activation capacity tests according to the present invention (herein referred to as "T cell assay") may preferably also be realised in transgenic mice, especially with suitably designed human MHC/HLA set up (e.g. having one or more human MHC/HLA molecules integrated in their genome).

Further preferred T cell assays are selected from T cell assays measuring phosphorylation/de-phosphorylation of components downstream of the T cell receptor, especially p56 lck, ITAMS of the TCR and the zeta chain, ZAP70, LAT, SLP-76, fyn, and lyn, T cell assays measuring intracellular Ca⁺⁺ concentration or activation of Ca⁺⁺-dependent proteins, T cell assays measuring formation of immunological synapses, T cell assays measuring release of effector molecules, especially perforin, granzymes or granulolysin or combinations of such T cell assays (see e.g. Current Protocols in Immunology, Current Protocols in Cell Biology).

WO 00/31542 presents methods for identifying antigens exclusively from tumor cells. The antigenic peptides are extracted from the MHC-peptide complexes located on the surface of hapten-modified malignant cells. Then haptenized peptides can be separated by hapten-specific affinity chromatography and sequenced. The embodiment of this invention is that peptides originally isolated from MHC molecules located on the surface of tumor cells have the property of stimulating T cells. Stimulation refers to T cell proliferation in response to the addition of cell extract, as well as production of cytokines such as INF-gamma, TNF, IL-2 and others. Based on the sequence of isolated peptide it is possible to identify the source of antigen.

Although the net result of this approach is also a T-cell epitope, this process substantially differs from the present invention: in the prior art natural MHC:peptide complexes are isolated from cellular systems, in the present invention purified, recombinant MHC molecules are used to isolate ligands from any source, cellular or synthetic, natural or artificial; in the prior art haptenization of the epitope is required for subsequent isolation by hapten-affinity chromatography, whereas the present invention is independent of such a step.

Tana et al. (1998) describe the approach for screening of antigenic peptides recognized by T cells from synthetic combinatorial peptide library designed on the base of the known binding motifs for set type of MHC molecules. Thus, the number of peptides to be screened is essentially reduced ($\sim 10^3$) comparing with a "comprehensive" library consisting of 20^9 all possible peptides. The peptides are combined in mixtures of nine peptides containing different amino acids in two fixed positions. Then the mixtures have been examined for eliciting T cell proliferative response. Thus, this approach is restricted for detection of amino acid residues appropriate for binding to the set MHC molecules and recognition by TCR receptors.

In Bitmansour et al. (2001) the matrix approach previously described by Kern et al., (1999, 2000) has been used for the identification of immunodominant CD4+ epitopes within CMV pp65 protein. A matrix of 24 peptide pools, each containing 12 pep-

- 12 -

tides (15-mers overlapping by 11 aa) (all 138 peptides) representing the whole pp65 protein was constructed. First, the peptide pools and later on the individual peptides have been checked for ability to provoke specific T cell response. The difference of this approach from others (Maeker, 2001; Tobery et al., 2001) is that it relies on the different techniques for monitoring T cell response, such as flow cytometric analysis (surface INF-gamma staining, CD4 and CD69 markers) and molecular methods (RT-PCR for TCR-V β content of CMV-specific CD4+ cells). The used technique, termed cytokine flow cytometry, allows detection of CMV-specific T-cells before either Ag induced proliferation or cell death, and therefore offers the possibility to determine the clonotypic content of CMV-specific T cells as it exists in vivo, unaltered by long-term in vitro culture. As a result, two pp65 epitopes (aa 489-503 and aa 509-523) contributing to presumably protective CMV-specific CD4+ response has been found in healthy CMV seropositive subjects.

Both, Tana et al. and Bitmansour et al., describe approaches which involve complex and thus only difficult to control biological, cellular assay, whereas the present invention represents a direct, biochemical isolation not comparable at all to the prior art, especially with respect to manageability and reproducibility: The use of "empty" MHC/HLA molecules according to the present invention makes all complex methods using binding events to MHC/HLA molecules being located on the surface of a living cell obsolete.

A method for isolating of T cell epitopes according to the present invention wherein a pool of ligands is contacted with ("empty") MHC/HLA molecules (and is not dependent on cellular systems) and then (preferably) the ligands which bind are tested in a T cell test with respect to their T cell activation capacity, is therefore a completely new approach, especially in comparison with the cell-dependent assays described. Also compared with "in silicio" methods, the present method provides "real world" utility with an easy and fast handling.

The performance and utility of the methods according to the present invention are demonstrated in the example section on a specific pathogen protein, pp65 of Cytomegalovirus (CMV). By

- 13 -

this example, the effectiveness and advantageousness of the present invention are shown and compared to prior art methods.

CMV, a betaherpesvirus, is the major cause of non-Epstein-Barr virus (EBV) infectious mononucleosis in the general population and an important pathogen in the immunocompromised host, including patients infected with the human immunodeficiency virus (HIV) suffering from acquired immunodeficiency syndrome (AIDS), neonates and transplant recipients (Drew et al, 1999). Depending on the population surveyed, the prevalence of CMV seropositivity in various regions ranges from 40-100%. As with other herpesviruses, primary CMV infection is followed by a persistent infection; re- or super-infection also occurs under certain circumstances (Britt, 1999), however mostly without pathologic consequences in the immunocompetent host because of pre-existing immunity (Plotkin et al, 1999). CMV establishes slow, persistent infections in humans. It is controlled, but never eliminated in the immunocompetent host. Among others (for example active expression of immunosuppressive genes, which interfere with antigen processing and presentation) two factors seem to contribute to this constant escape from immunosurveillance: replication in immunoprivileged sites like salivary gland epithelia on the one hand and the formation of a latent reservoir in CD33+ monocytes. These macrophage precursors do not support replication of CMV, which is therefore arrested until cells differentiate into macrophages. Only after differentiation the cellular environment seems to become permissive for lytic infection. 90% of primary infections in immunocompetent individuals go clinically unrecognized (Nichols et al, 2000) and long-term immunity develops, which controls viral persistence and - albeit not being sterilizing - is protective.

A high risk of infection/reactivation exists for the following groups, who are highly jeopardized to develop CMV disease:

- a) Fetuses from seronegative pregnant women
- b) seronegative transplant patients, because it is difficult to find CMV negative grafts,
- c) seropositive transplant and HIV patients, because their induced or acquired immunodeficiency allows for reactivation of CMV from latency.

Congenital CMV infection may lead to deafness and more important is as frequent a cause of mental retardation as the common genetic syndromes, trisomy 21 and fragile X chromosome. From the study of Fowler KB, Stagno S, Pass RF, et al. (Fowler et al, 1992) one can extrapolate, that about 9000 European and 8000 American infants (representing 1 out of 500 infants born in the United states) are harmed each year as a result of intrauterine CMV infection, of which only 10% are clinically apparent at birth. Although congenital CMV infection is largely silent at birth, its cumulative effect is large in terms of clinical sequelae and public health impact (Plotkin et al, 1999). The factor, which is most closely associated with poor outcome, is a primary maternal infection during gestation, which is not easily to be prevented in seronegative mothers, because the virus spreads easily by close person-to-person contact and is shed in high amounts by seropositive toddlers (Field et al, 1999). In the immunocompromised adult, disease is most frequently seen in solid organ transplant (SOT) patients and allogeneic hematopoietic stem cell transplant (HCT) patients as well as in HIV infected individuals. In allogeneic HCT patients severe CMV disease still occurs in about 15% of the patients, with a mortality of about 50%. The risk factors associated with CMV disease in HCT are a CMV positive graft donor and an uninfected graft recipient, concomitant bacterial infections, fulminant hepatitis and graft-versus-host disease. For SOT such as those of kidney, liver, heart, lung or the pancreas, CMV disease is associated with decreased graft and patient survival. CMV causes a variety of infectious diseases syndromes itself. The consequences of CMV disease are similar in all transplant patients, although specific organ involvement frequently corresponds to the organ transplanted. In general, liver, pancreas, lung, intestinal, and heart transplant recipients have a greater incidence of CMV disease than kidney transplant recipients. Symptomatic infections occur in approximately 39-41% of heart-lung, 9-35% of heart, 22-29% of liver and pancreas, and 8-32% of renal transplant recipients not receiving antiviral prophylaxis. Moreover is CMV infection associated with an augmented immunosuppressed state, which may explain the frequent opportunistic superinfections in transplant patients. Furthermore it seems to

be involved in allograft dysfunction and indirectly in decreased patient survival as well as increased costs and longer hospital lengths of stay (Sia et al, 2000). To the latter three points CMV's suspected involvement on the one hand in accelerated coronary arteriosclerosis found in patients with CMV infection after heart transplantation (Van Son et al, 1999 and Field et al, 1999) and on the other hand in the enhanced risk of Epstein-Barr Virus (EBV) related posttransplantation lymphoproliferative disease in CMV/EBV doublepositive transplant patients (Sia et al, 2000) might contribute. CMV retinitis has been one of the most common disease manifestations in AIDS patients and has been reported to occur in about 30% of patients, threatening those patients with blindness. Recent evidence indicates that the loss of CD4+ lymphocytes correlates with the development of CMV disease in AIDS patients and therefore the introduction of HIV protease inhibitors and highly active antiretroviral therapy (HAART), the incidence of CMV retinitis has declined significantly (Field et al, 1999). However CMV disease can still occur within 4 months of HAART initiation despite adequate suppression of HIV replication. Furthermore antiretroviral therapy failure is a problem on the rise (Nichols et al, 2000).

In the last decade, considerable progress has been made in the use of antiviral chemotherapy to prevent and treat CMV infection. At present there are five drugs that have been approved in the US for the treatment of CMV: Ganciclovir (Cytovene, when used as intravenous and oral formulation or Vitrasert, when used as intravitreal implant formulation), Foscarnet (Foscavir), Cidofovir (Vistide) and Fomivirsen (Vitracene). The triphosphate equivalents of Ganciclovir and Cidofovir (requiring phosphorylation by viral and cellular enzymes in case of the former and cellular enzymes only in case of the latter) are competitive inhibitors of deoxyribonucleotidetriphosphates for the viral DNA polymerase, while the pyrophosphate analogue Foscarnet blocks the pyrophosphate-binding site of this enzyme. Fomivirsen is the first antisense designed oligonucleotide to gain FDA approval. However the mechanism of action of Fomivirsen has never been fully confirmed and may have many components besides antisense (Field et al, 1999). Because of inherent high toxicities, high incidence of late CMV disease and high costs of universal pro-

- 16 -

phylaxis, a preemptive strategy of antiviral therapy emerged that is based on the selective treatment of patients with a high risk for developing CMV disease based on early detection of re-activated CMV post-HCT (Zaia et al, 2000). In SOT patients prophylactic strategies are particularly attractive in the setting of a graft donor CMV carrier and an uninfected graft recipient. Valganciclovir, the valine ester of the active drug ganciclovir with markedly increased oral bioavailability, given for three months reduced the incidence of CMV disease in kidney transplant patients markedly. Moreover there was also a significant reduction in the proportion of patients with biopsy-proven rejection at six months after transplantation. However, long-term prophylaxis especially with available oral agents, which make this form of treatment more convenient for the patient than conventional intravenous formulation, may also increase the incidence of drug resistance (Nichols et al, 2000). Multidrug resistance to Ganciclovir, Cidofovir and Foscarnet -all due to changes in the viral DNA polymerase- has already been observed (Field et al, 1999). Fomivirsen seems to be effective against such mutants (Nichols et al, 2000) but because of its probable antiviral activity besides its antisense mode of action (Field et al, 1999), Fomivirsen resistant strains seem likely to arise.

Therefore a focus was set on the (re)constitution of CMV specific immunity after transplantation, which might also prevent damages to babies of seronegative mothers. This is related to the question, how the immunocompetent host is controlling CMV infection, so as to learn, what is necessary to be restored in the non-immunocompetent host. Aggregate data argue that CMV antibodies are partly protective through reduction of viremia (dissemination of cell-free virus), but are little effective in HCT patients, where the virus persists cell-associated within the monocyte population, a reservoir, which cannot be destroyed by the use of antibodies (Plotkin et al, 1999). This argues strongly for the development of vaccines, which restore CD8+ and CD4+ T cell responses. The protective function of CMV specific alphabeta/T cells could definitively be established by adoptive transfer of these cells early after transplantation. In recipients of adoptively transferred CD8+ CMV-specific cytotoxic T lymphocytes (CTL) in a phase I study, cytolytic responses equi-

- 17 -

valent to those in the immunocompetent marrow donor were achieved immediately after the fourth infusion. However they declined over the ensuing several weeks in the subset of patients who failed to recover endogenous CD4+ CMV-specific T helper responses, which underlines the importance of generating both CD8+ and CD4+ T cell responses (Zaia et al, 2000). The potential importance of CD4+ T cells in CMV control is also suggested by the very high frequencies of specific CD4+ memory cells (about 2.0% of total CD4+ T cells) in normal CMV seropositive individuals (Waldrop et al, 1998). The close association between the degree of CD4+ T cell deficiency and CMV disease in HIV infection is also thought to be consistent with a crucial role for CD4+ T cells in control of CMV reactivation (Komanduri et al, 1998). Five general types of CMV-related vaccines have been described: attenuated live virus vaccines, recombinant live virus vaccines, DNA vaccines, whole protein vaccines, and peptide vaccines. An attenuated live virus vaccine, the "Towne strain", was developed and tested in the 1970s both in renal transplant patients and women in childbearing years. Although this vaccine was shown to be immunogenic, the use of a live virus in the transplant population presents a potential risk. To a lesser extent this also holds true for recombinant poxviruses with limited potential for replication in humans (avipox). Approaches, that show promise in the animal model include introduction of DNA vectors encoding immunogenic proteins as a means to elicit CTL. Refinement of DNA vaccine technology, including the use of minimal cytotoxic and helper T cell epitopes as immunogens may result in even more efficient vaccines (Zaia et al, 2000).

Taken together the negative side effects and high costs of anti-viral drugs, the limited application range/ success of above mentioned approaches for a CMV vaccine and the good efficacy of adoptive transfer of T cell clones highlights the importance of finding new MHC class I and II restricted T cell epitopes. These epitopes should be presented by the most prevalent HLA molecules and therefore confer protection to the majority of the population irrespective of the individual HLA setting. CMV is one of the viruses with the highest protein-coding capacity known, with 170-200 open reading frames (Reddehase, 2000). Analysis of the T cell responses of asymptomatic donors, obviously controlling the

- 18 -

virus successfully, however revealed the immunodominance of CMV pp65, pp150, IE-1 and gB (Zaia et al, 2000). This is probably due to the expression of several immunosuppressive genes, which interfere with the formation and egress of peptide loaded MHC class I and II molecules. Remarkably, the downregulation of MHC class I molecules seems to interfere little with recognition of CMV-infected cells by CMV pp65 or pp150-specific CTL. Very early after viral entry, during which these 2 structural proteins are delivered into the cytoplasm, this can be explained by the fact that these structural proteins can be processed and presented before the expression of immunosuppressive viral genes. CMV pp65 or pp150 specific CTL however, recognize infected cells also efficiently at later stages, when MHC levels are already reduced. This seems to be due to a host counter evasion strategy. Upon CMV infection a cellular gene is induced, which binds to a receptor on T cells and facilitates T cell activation even when the target cell expresses only a low peptide/MHC density (Zaia et al, 2000). On the other hand, there are indications, that pp65 interferes with the presentation of IE-1, the activator of all ensuing immunosuppressive gene products (Reddehase, 2000), which could explain the immunodominance of the pp65 antigen. Thus, T cells specific for pp65 are able to lyse virus infected cells at all stages of the replication cycle and may be essential for eliminating infected cells in vivo (Zaia et al, 2000).

According to a further aspect, the present invention also provides T cell epitopes identifiable by a method according to the present invention, said T cell epitopes being selected from the group consisting of polypeptides comprising the sequence KM-QVIGDQYVK, FTWPPWQAGI, AMAGASTSA, SDNEIHNPV, KYQEFFWDA or combinations thereof.

The present invention also provides HLA A0201 binding epitopes with T cell activating capacity identifiable by a method according to the present invention using HLA A0201 molecules as MHC/HLA molecules, said HLA A0201 binding epitopes being selected from the group consisting of polypeptides comprising the sequence RLLQTGIHV, VIGDQYVKV, YLESCFEDV or combinations thereof. Although these sequences have been known to bind to HLA A0201, their useability to activate T cells is provided with the

- 19 -

present invention.

The present invention further provides the use of a peptide comprising the sequence RPHERNNGFTV for preparing a composition for activating T cells in an individual being B7-negative.

Additionally, the present invention provides the use of a peptide comprising the sequence DDVWTSGSDSDE for preparing a composition for activating T cells in an individual being B35-negative.

Moreover, the present invention also provides the use of a peptide comprising the sequence TPRVTGGGAM for preparing a composition for activating T cells in an individual being B7-negative.

Although these sequences have been described to have a specific HLA restriction (RPHERNNGFTV: B7, DDVWTSGSDSDE: B35, TPRVTGGGAM: B7), it was surprising that these sequences have a specificity being different from the known restriction (RPHERNNGFTV: non-B7, DDVWTSGSDSDE: non-B35, TPRVTGGGAM: non-B7). This enables an expansion of the usefulness of these sequences, e.g. by making these sequences available as suitable vaccines for individuals expressing other HLAs than the ones described.

The present invention further provides peptides binding to class II HLA molecules selected from peptide nos. 55-64, 109, 383, 384, 421, 449-454, 469 and 470 according to table 3 of the example section.

Preferably, the epitopes or peptides according to the present invention further comprises 1 to 30, preferably 2 to 10, especially 2 to 6, naturally occurring amino acid residues at the N-terminus, the C-terminus or at the N- and C-terminus. For the purposes of the present invention the term "naturally occurring" amino acid residue relates to amino acid residues which are present in the naturally occurring protein at the specific position, relative to the epitope or peptide. For example, for the "AMAGASTSA" epitope, the naturally occurring amino acid residue at the N-terminus is Gly; the three naturally occurring amino acid residues at the C-terminus are Gly-Arg-Lys. A "non-natur-

- 20 -

ally occurring" amino acid residue is therefore any amino acid residue being different as the amino acid residue at the specific position relative to the epitope or peptide.

According to a preferred embodiment of the present invention, the present epitopes or peptides further comprise non-naturally occurring amino acid(s), preferably 1 to 1000, more preferred 2 to 100, especially 2 to 20 non-naturally occurring amino acid residues, especially at the N-terminus, the C-terminus or at the N- and C-terminus. Also combinations of non-naturally and naturally occurring amino acid residues are possible under this specific preferred embodiment. The present epitope may also contain modified amino acids (i.e. amino acid residues being different from the 20 "classical" amino acids, such as D-amino acids or S-S bindings of Cys) as additional amino acid residues or in replacement of a naturally occurring amino acid residue.

It is clear that also epitopes or peptides derived from the present epitopes or peptides by amino acid exchanges improving, conserving or at least not significantly impeding the T cell activating capability of the epitopes are covered by the epitopes or peptides according to the present invention. Therefore, the present epitopes or peptides also cover epitopes or peptides, which do not contain the original sequence as derived from CMV pp65, but trigger the same or preferably an improved T cell response. These epitopes are referred to as "heteroclitic". These include any epitope, which can trigger the same T cells as the original epitope and has preferably a more potent activation capacity of T cells preferably in vivo or also in vitro.

Heteroclitic epitopes can be obtained by rational design i.e. taking into account the contribution of individual residues to binding to MHC/HLA as for instance described by Ramensee et al. 1999 or Sturniolo et al. 1999, combined with a systematic exchange of residues potentially interacting with the TCR and testing the resulting sequences with T cells directed against the original epitope. Such a design is possible for a skilled man in the art without much experimentation.

Another possibility includes the screening of peptide libraries

- 21 -

with T cells directed against the original epitope. A preferred way is the positional scanning of synthetic peptide libraries. Such approaches have been described in detail for instance by Blake et al 1996 and Hemmer et al. 1999 and the references given therein.

As an alternative to epitopes represented by the cognate CMV pp65 derived amino acid sequence or heteroclitic epitopes, also substances mimicking these epitopes e.g. "peptidemimetica" or "retro-inverso-peptides" can be applied.

Another aspect of the design of improved epitopes is their formulation or modification with substances increasing their capacity to stimulate T cells. These include T helper cell epitopes, lipids or liposomes or preferred modifications as described in WO 01/78767.

Another way to increase the T cell stimulating capacity of epitopes is their formulation with immune stimulating substances for instance cytokines or chemokines like interleukin-2, -7, -12, -18, class I and II interferons (IFN), especially IFN- γ , GM-CSF, TNF-alpha, flt3-ligand and others.

According to a further aspect, the present invention is drawn to the use of an epitope or peptide according to the present invention for the preparation of a HLA restricted vaccine for treating or preventing cytomegalovirus (CMV) infections.

The invention also encompasses the use of an epitope comprising the sequence KMQVIGDQYV, FTWPPWQAGI, AMAGASTSA, SDNEIHNPV and/or KYQEFFWDA for the preparation of a vaccine for treating or preventing cytomegalovirus (CMV) infections.

Consequently, the present invention also encompasses a vaccine for treating or preventing cytomegalovirus (CMV) infections comprising an epitope comprising the sequence KMQVIGDQYV, FTWPPWQAGI, AMAGASTSA, SDNEIHNPV and/or KYQEFFWDA. Furthermore, also a HLA specific vaccine for treating or preventing cytomegalovirus (CMV) infections comprising the epitopes or peptides according to the present invention is an aspect of the present

invention. The application of the corresponding nucleic acids encoding the peptides according to the present invention, e.g. as DNA vaccine, also falls under the scope of the present invention, at least as equivalent to the claimed peptide vaccines.

Parker et al. (1994) present the method of prediction of peptide binding to MHC class I molecules based on the experimental data of binding of individual peptides. The peptide binding ability was assessed indirectly by monitoring the ability of peptides to promote incorporation of beta2-microglobulin (β 2m) into HLA-A2/ β 2m/ peptide heterodimeric complexes. The experimental data (measured rates of dissociation of β 2m) allows to create the value of corresponding coefficients used then to calculate theoretical binding stability for any nonapeptides in the complex with HLA-A2 molecules. In this study it has been shown for the first time that CMV pp65 derived peptide sequence RLLQTGIHV can stabilize HLA-A2/ β 2m/ peptide complex.

Morgan et al. (1998) describe the method to determine the binding affinity of peptides to purified HLA molecules also based on the indirect measurement of the incorporation of β 2m into the HLA/peptide complex. In this study the in vitro binding of RLLQTGIHV peptide has been demonstrated and its relative binding affinity to HLA-A2 molecule (association and dissociation constants) was measured.

Although binding to HLA-A2 is demonstrated by Parker et al. and Morgan et al., the final proof that this ligand also represents a true T-cell epitope is lacking, whereas in the present invention this is demonstrated by means of interferon-gamma ELISPOT proofing, that RLLQTGIHV can not only bind to HLA-A2, but that it can only induce functional T-cells. This is not a trivial finding: it is well known in the field that many ligands binding with high affinity do not represent T-cell targets, as they are for instance not or not efficiently generated through the aforementioned "antigen processing and presentation pathway. Therefore, HLA A0201 binding activity of RLLQTGIHV was not only surprising but may also be selectively used in vaccines specifically designed e.g. for a certain allele population.

Preferably, such a vaccine according to the present invention further comprises an immunomodulating substance, preferably selected from the group consisting of polycationic substances, especially polycationic polypeptides, immunomodulating nucleic acids, especially deoxyinosine and/or deoxyuracile containing oligodeoxynucleotides, or mixtures thereof.

Preferably the vaccine further comprises a polycationic polymer, preferably a polycationic peptide, especially polyarginine, polylysine or an antimicrobial peptide.

The polycationic compound(s) to be used according to the present invention may be any polycationic compound which shows the characteristic effect according to the WO 97/30721. Preferred polycationic compounds are selected from basic polypeptides, organic polycations, basic polyaminoacids or mixtures thereof. These polyaminoacids should have a chain length of at least 4 amino acid residues. Especially preferred are substances containing peptidic bounds, like polylysine, polyarginine and polypeptides containing more than 20%, especially more than 50% of basic amino acids in a range of more than 8, especially more than 20, amino acid residues or mixtures thereof. Other preferred polycations and their pharmaceutical compositions are described in WO 97/30721 (e.g. polyethyleneimine) and WO 99/38528. Preferably these polypeptides contain between 20 and 500 amino acid residues, especially between 30 and 200 residues.

These polycationic compounds may be produced chemically or recombinantly or may be derived from natural sources.

Cationic (poly)peptides may also be polycationic anti-bacterial microbial peptides. These (poly)peptides may be of prokaryotic or animal or plant origin or may be produced chemically or recombinantly. Peptides may also belong to the class of defensins. Such host defense peptides or defensins are also a preferred form of the polycationic polymer according to the present invention. Generally, a compound allowing as an end product activation (or down-regulation) of the adaptive immune system, preferably mediated by APCs (including dendritic cells) is used as polycationic polymer.

Especially preferred for use as polycationic substance in the present invention are cathelicidin derived antimicrobial peptides or derivatives thereof (A 1416/2000, incorporated herein by reference), especially antimicrobial peptides derived from mammal cathelicidin, preferably from human, bovine or mouse, or neuroactive compounds, such as (human) growth hormone (as described e.g. in WO01/24822).

Polycationic compounds derived from natural sources include HIV-REV or HIV-TAT (derived cationic peptides, antennapedia peptides, chitosan or other derivatives of chitin) or other peptides derived from these peptides or proteins by biochemical or recombinant production. Other preferred polycationic compounds are cathelin or related or derived substances from cathelin, especially mouse, bovine or especially human cathelins and/or cathelicidins. Related or derived cathelin substances contain the whole or parts of the cathelin sequence with at least 15-20 amino acid residues. Derivations may include the substitution or modification of the natural amino acids by amino acids which are not among the 20 standard amino acids. Moreover, further cationic residues may be introduced into such cathelin molecules. These cathelin molecules are preferred to be combined with the antigen/vaccine composition according to the present invention. However, these cathelin molecules surprisingly have turned out to be also effective as an adjuvant for a antigen without the addition of further adjuvants. It is therefore possible to use such cathelin molecules as efficient adjuvants in vaccine formulations with or without further immunactivating substances.

Another preferred polycationic substance to be used according to the present invention is a synthetic peptide containing at least 2 KLK-motifs separated by a linker of 3 to 7 hydrophobic amino acids, especially L (e.g. KLKL₅KLK; PCT/EP01/12041, incorporated herein by reference).

The immunomodulating (or:immunogenic) nucleic acids to be used according to the present invention can be of synthetic, prokaryotic and eukaryotic origin. In the case of eukaryotic origin, DNA should be derived from, based on the phylogenetic tree, less

developed species (e.g. insects, but also others). In a preferred embodiment of the invention the immunogenic oligodeoxynucleotide (ODN) is a synthetically produced DNA-molecule or mixtures of such molecules. Derivates or modifications of ODNs such as thiophosphate substituted analogues (thiophosphate residues substitute for phosphate) as for example described in US patents US 5,723,335 and US 5,663,153, and other derivatives and modifications, which preferably stabilize the immunostimulatory composition(s) but do not change their immunological properties, are also included. A preferred sequence motif is a six base DNA motif containing an (unmethylated) CpG dinucleotide flanked by two 5' purines and two 3' pyrimidines (5'-Pur-Pur-C-G-Pyr-Pyr-3'). The CpG motifs contained in the ODNs according to the present invention are more common in microbial than higher vertebrate DNA and display differences in the pattern of methylation. Surprisingly, sequences stimulating mouse APCs are not very efficient for human cells. Preferred palindromic or non-palindromic ODNs to be used according to the present invention are disclosed e.g. in Austrian Patent applications A 1973/2000, A 805/2001, EP 0 468 520 A2, WO 96/02555, WO 98/16247, WO 98/18810, WO 98/37919, WO 98/40100, WO 98/52581, WO 98/52962, WO 99/51259 and WO 99/56755 all incorporated herein by reference. Apart from stimulating the immune system certain ODNs are neutralizing some immune responses. These sequences are also included in the current invention, for example for applications for the treatment of autoimmune diseases. ODNs/DNAs may be produced chemically or recombinantly or may be derived from natural sources. Preferred natural sources are insects.

Alternatively, also nucleic acids based on inosine and cytidine (as e.g. described in the PCT/EP01/06437) or deoxynucleic acids containing deoxy-inosine and/or deoxyuridine residues (described in the Austrian patent applications A 1973/2000 and A 805/2001, incorporated herein by reference) may preferably be used as immunostimulatory nucleic acids for the present invention.

Of course, also mixtures of different immunogenic nucleic acids may be used according to the present invention.

Preferably, the present vaccine further comprises a pharmaceut-

ically acceptable carrier.

According to a further preferred embodiment, the present vaccine comprises an epitope or peptide which is provided in a form selected from peptides, peptide analogues, proteins, naked DNA, RNA, viral vectors, virus-like particles, recombinant/chimeric viruses, recombinant bacteria or dendritic cells pulsed with protein/peptide/RNA or transfected with DNA comprising the epitopes or peptides.

According to a further aspect, the present invention is drawn to T cells, a T cell clone or a population (preparation) of T cells specifically recognizing any epitope or peptide according to the present invention, especially a CMV epitope as described above. A preferred application of such T cells is their expansion in vitro and use for therapy of patients e.g. by adoptive transfer. Therefore, the present invention also provides the use of T cells, a T cell clone or a population (preparation) of T cells for the preparation of a composition for the therapy of CMV patients.

Such T cells (clones or lines) according to the present invention, specifically those recognizing the aforementioned CMV peptides are also useful for identification of heteroclitic epitopes, which are distinct from the originally identified epitopes but trigger the same T cells.

Such cells, compositions or vaccines according to the present invention are administered to the individuals in an effective amount.

The invention will be explained in more detail by way of the following examples and drawing figures, to which, however it is not limited.

Fig.1 shows peptide binding affinities to soluble DR4 molecules;

Fig.2 shows identification of peptides capable of binding to empty DR4 molecules (A. Purification of HLA-peptide complexes; B. MS analysis of bound peptides);

- 27 -

Fig. 3 shows the binding of high affinity peptide in the presence of the excess of low affinity peptide to DR4 molecules;

Fig. 4 shows the binding of the individual peptides and peptide mixtures to DR4 molecules;

Fig.5 shows an ELISPOT with (5a) isolated CD4+ T cells and DR0401 molecules (costimulation with anti CD28) and with (5b) isolated CD8+ T cells and HLA A0201 molecules (costimulation with anti CD28 mab);

Fig.6 shows the CMV pp65 peptide pool array;

Fig.7 shows (7a) the binding of peptide pools derived from pp65 protein to DR4 molecules and (7b) binding of single pp65 peptides to DR4 molecules;

Fig.8 shows the 1st screen with peptide mixtures (containing 21 peptides each, Donor #10736 HLA A2/3, HLA B15/35; 8a) and the 1st screen with peptide mixtures (containing 21 peptides each, Donor #10687 HLA A2/11, HLA B7/13; 8b);

Fig.9 shows the 2nd screen with single peptides (Donor #10736 HLA A2/3, HLA B15/35);

Fig. 10 shows PBMC from subject 10788 applied for IFN- γ ELISPOT with CMVpp65 15mers 57, 59 and controls (med: no peptide, HIV: irrelevant HIV-derived peptide, ConA: polyclonal stimulation;

Fig. 11 shows confirmation of simultaneous CD4+ and CD8+ T-cell responses against CMVpp65 15mers 469, 470 by intracellular IFN- γ staining; and

Figure 12 shows mapping of DRB1*0401 epitopes with overlapping 15 mers in transgenic mice using IFN- γ ELISPOT assay: One week later after the last vaccination, spleens were removed and cells were activated ex vivo with relevant peptides (no. 1500-1505), overlapping 15mers representing these longer peptides and irrelevant influenza hemagglutinin derived peptide (no.1171) to de-

- 28 -

termine IFN- γ -producing specific cells (medium control is subtracted).

Examples :

General description of the examples:

The present examples show the performance of the present invention on a specific pathogen protein, pp65 of CMV.

In the first part the method according to the present invention was applied, which is based on the use of "empty HLA molecules". These molecules were incubated with mixtures of potential CMV derived peptide ligands, screening for specific binding events. The complexes formed in this way were isolated and used for the identification of the specifically bound ligands. The possibility to use highly complex mixtures allows a very quick identification of the few binders out of hundreds or even thousands of potential ligands. This is demonstrated by using HLA-DRB1*0401 and pools of overlapping 15-mers.

However, the same process can be applied for class I molecules and peptides of appropriate length i.e. 8 to 11-mers. The ligand-pools can be synthetic overlapping peptides. Another possibility is to digest the antigen in question enzymatically or non-enzymatically. The latter achieved by alkali-hydrolysis generates all potential degradation products and has been successfully used to identify T cell epitopes (Gavin 1993). Enzymatic digestions can be done with proteases. One rational way would further be to use proteases involved in the natural antigen-processing pathway like the proteasome for class I restricted epitopes (Heemels 1995) or cathepsins for class II restricted epitopes (Villadangos 2000). Ligand pools could also be composed of naturally occurring ligands obtained for instance by lysis of or elution from cells carrying the respective epitope. In this regard it is important to note that also non-peptide ligands like for instance glycolipids can be applied. It is known that nonclassical class I molecules, which can be encoded by the MHC (e.g. HLA-G, HLA-E, MICA, MICB) or outside the MHC (e.g. CD1 family) can present various non-peptide ligands to lymphocytes

(Kronenberg 1999). Use of recombinant "empty" nonclassical class I molecules would allow binding reactions and identification of binders in similar manner as described here.

After rapid identification of ligands capable of binding to HLA molecules the process according to the present invention also offers ways to characterize directly specific T cell responses against these binders. One possibility is to directly use the isolated HLA:ligand complex in a so called "synthetic T cell assay". The latter involves antigen-specific re-stimulation of T cells by the HLA:ligand complex together with a second signal providing co-stimulation like activation of CD28 by an activating antibody. This assay can be done in an ELIspot readout as demonstrated in Example II.

Here, it was chosen to synthesize the CMV pp65 as a series of overlapping 15 mer peptides, each peptide overlapping its precursor by 14 out of 15 aa. The peptides were supplied as pools of 21 single peptides, which were constructed that way that each single peptide occurs in exactly 2 pools. Array of the peptide pools in matrix form allows identification of single peptides as the crossover points of row- and column mixtures (Fig. 6). As donors 10 healthy CMV seropositive blood donors, expressing the MHC class I molecule A2, were chosen. This MHC preference was inferred because it is known, that the pp65 antigen is especially well recognized by donors carrying this MHC molecule (Saulquin et al, 2000; Kern et al, 2000).

In parallel the same peptides (pools) were screened by interferon-gamma (IFN- γ) ELIspot assays using peripheral blood mononuclear cells (PBMC) from CMV seropositive individuals (binding to soluble recombinant HLA-class II molecules in an SDS-stability assay). Both approaches yielded epitopes, and although not biased by the choice of HLA (class I HLA-A2 for the ex vivo T cell assay approach, class II HLA-DR4 for the peptide binding approach) showed some degree of overlap.

MATERIALS & METHODS

Peptides

547 fifteen amino acid residue (15mer) peptides overlapping by 14 aa representing the complete sequence of the CMV pp65 antigen, were synthesized on a Syro II synthesizer (MultisynTech, Witten, Germany). 288 peptides were made in parallel using standard F-moc chemistry.

Each peptide was solubilized in 100 % DMSO at ~10 mg/ml. Stocks of 42 peptide pools derived from pp65 were made in 100 % DMSO at a final concentration of 0.45 mg/ml for each peptide.

The other peptides were synthesized using standard F-moc chemistry either on the Syro II synthesizer or a ABI 433A synthesizer (Applied Biosystems, Weiterstadt, Germany) and purified by RP-HPLC (Biocut 700E, Applied Biosystems, Langen, Germany) using a C18 column (either ODS ACU from YMC or 218TP from Vydac). Purity and identity of each peptide were characterized by MALDI-TOF on a Reflex III mass-spektrometer (Bruker, Bremen, Germany).

Peptide binding assay

Soluble HLA class I A*0201 and HLA class II DRA1*0101/DRB1*0101/Ii, DRA1*0101/DRB1*0401/Ii and DRA1*0101/DRB1*0404/Ii molecules were expressed in SC-2 cells and purified as described in Aichinger et al., 1997.

In peptide binding reactions HLA molecules were used in a concentration of 0.5 μ M, and each single peptide was added in 10-fold molar excess (5 μ M) if not mentioned differently. The concentration of DMSO in the binding reaction did not exceed 4 %. The reaction was performed in PBS buffer (pH 7.4) at room temperature for 48 hours in the presence of a protease inhibitor cocktail (Roche) and 0.1 % octyl-b-D-glucopyranoside (Sigma).

Peptide binding was evaluated in an SDS-stability assay (Gorga et al., 1987): trimeric HLA class II ab-peptide complexes are resistant to SDS and consequently appear as ~60 kDa band in SDS-

- 31 -

PAGE Western blot analysis. Individual HLA class II α and β chains not stabilized by a peptide binding with intermediate to high affinity migrate as ~35 kDa and ~30 kDa bands, respectively.

Briefly, HLA-peptide complexes were treated with 1 % SDS at room temperature and resolved by SDS-PAGE run with 20 mA for approximately 2.5 hours at room temperature. Protein was transferred onto PVDF membrane by electroblotting, and stained with anti- α -chain TAL.1B5 α or/and β -chain MEM136 antibodies. For detection of Western-blot signals ECL solutions (Amersham) were used.

The binding affinities to DRB1*0701 and DRB1*1101 were tested by a peptide-competition assay (Reay et al., 1992). Briefly, binding of the biotinylated CLIP peptide (reference peptide) has been used for monitoring of HLA:peptide complex formation. A testing peptide added to the binding reaction at an equimolar concentration to CLIP peptide could compete out CLIP when its affinity is higher or inhibit binding for 50 % if its affinity is equal to affinity of CLIP. In the case of lower affinity peptides they should be added in excess to the reference peptide to compete for occupancy of HLA binding groove. The values of the concentration of competitor peptides required for 50 % inhibition of reference peptide binding (IC_{50}) can be used for evaluation of peptide binding affinities. Alternatively, comparing of the amount of reference peptide bound to HLA molecules in the presence or absence of competitor peptide one can determine the binding activity of the peptide of interest.

In the present peptide-competition assay conditions of peptide binding were similar to described above. Soluble HLA molecules were used in the concentration of 0.5 μ M and biotinylated CLIP was added to all samples in the final concentration of 2 μ M. Competitor peptides were added in three different concentrations: 0.25, 5 and 100 μ M. Binding reaction was performed in PBS buffer (pH 7.4) for 18 hours at 37°C. Amount of biotinylated CLIPm, associated with soluble HLA molecules, was determined by ELISA. Briefly, MaxiSorb 96-well plates (Nunc, Denmark) were coated with mouse anti- $\alpha\beta$ antibody L243 (purified from ATCC HB-55) by overnight incubation with 50 μ l of 10 μ g/ml dilution in

- 32 -

PBS at 4°C. Non-specific binding to the plastic was blocked by incubation with T-PBS containing 3 % of BSA for 2 hours at 37°C and binding reactions were then "captured" for 2 hours at room temperature. After extensive washing of the plates with T-PBS, HLA-associated peptide complexes were detected colorimetrically using alkaline phosphatase-streptavidin conjugate (Dako) and Sigma 104 phosphatase substrate (Sigma Diagnostics, USA). The optical density at 405 nm was measured on a microplate reader SUNRISE (Tecan).

Identification of peptides from HLA-peptide complexes

After peptide binding reaction with soluble HLA molecules, HLA-peptide complexes were separated from free peptides by gel-filtration chromatography on Superdex-200 column (ÄKTA design, Amersham Pharmacia Biotech). HLA-containing fractions were collected, and bound peptides were reconstituted from the complexes by adding TFA to the final concentration of 1 %. Peptides were desalted by Ziptip purification and analyzed by MALDI-TOF mass-spectrometry.

Synthetic Elispot Assays

96well filtration plates from Millipore (catalog-number: MAHA S4510) were coated with a mixture of 1 µg/well anti human IFN-γ mab B140 from Bender Med Systems and 0,5 µg/well QIAexpress Penta.His antibody from Qiagen over night at 4°C. As a positive control for viability and cytokine production of CD8+ T cells some wells were coated with anti CD3 antibody, clone MEM-57 from V. Horejsi, Institute of Molecular Genetics, Prag.

Plates were washed 2 times with PBS (from GIBCOBRL, catalognumber 14190-094) and blocked with the following ELISPOT medium: RPMI 1640 from GIBCOBRL (catalog number: 31870-025) supplemented with 1 mM sodium pyruvate from GIBCOBRL (catalog number: 11360-039), 2 mM L-glutamine from GIBCOBRL (catalog number: 25030-024), 0,1 mM non-essential amino acids from GIBCOBRL (catalog number: 11140-035), 50 µg/ml gentamycin from GIBCOBRL (catalog number: 15710-049), 50 µM 2-mercaptoethanol from GIBCOBRL (catalog number: 31350-010) and 10% human serum type AB from

- 33 -

BioWhittaker (catalog number: 14-490E) for 30 min at 37°C .

After removal of the blocking medium wells were incubated with soluble HLA DRB1*0401 loaded with either peptide 1242 derived from M. tuberculosis, peptide 1171 derived from Influenza Hemagglutinin (HA-pep: aa 306-318 or for a negative control peptide 84 derived from Hepatitis C virus (HCV-pep: NS3 aa1248-1261), diluted in ELISPOT medium to 100 µg/ml (10µg/well), for 5 hours at 37°C (Fig. 5a). For fig. 5b soluble HLA A*0201 molecules loaded either with a peptide derived from the EBV antigen BMLF1 aa280-88, peptide 21 derived from Influenza Matrix protein aa 58-66 or for a negative control peptide 90 derived from HIV reverse transcriptase aa 476-484 were used. Loading of molecules was essentially done as described in the paragraph "Peptide binding assay".

Mononuclear cells from the peripheral blood (PBMC) of healthy, BCG-vaccinated HIV and HCV negative donors with matching HLA phenotype were isolated on Lymphoprep (from Nycomed Pharma AS, Oslo, Norway) using Leuco Sep tubes (from Greiner), washed 3x with PBS (from GIBCOBRL, catalognumber 14190-094).

From the PBMC either the CD4+ T cells (Fig. 5a) or the CD8+ T cells (Fig. 5b) were isolated using MACS technique (Miltenyi, Germany) according to the manufacturer's instructions. The isolated T cells were resuspended in ELISPOT medium containing 10 µg/ml anti CD28 monoclonal antibody (clone 37407.111, mab 342 from R&D systems) at a concentration of 1 Mio/ml.

The solution containing the soluble HLA (sHLA) molecules was discarded and the isolated T cells were seeded into the wells. The respective samples were resupplemented with the according peptides at 5 µM (Fig. 5a) or 10 µg/ml (Fig. 5b).

Cells were cocultivated with the sHLA molecules for 20 hrs. Assays were arrested by shaking off the contents and washing 6x with wash buffer (PBS; 0,1% Tween 20 from SIGMA). Next 100 µl of a 1:10000 dilution of the biotinylated anti human IFN-γ mab (B308-BT2 from BMS), which corresponds to 0,015 µg/well, was added for an incubation of 2 hrs at 37°C or alternatively for over

- 34 -

night at 4°C. After washing, Streptavidin-ALP from DAKO (catalog number: D0396) was added at 1,2 µg/ml for 1 hr at 37°C. The assay was developed by addition of 100 µl/well BCIP/NBT alkaline phosphatase substrate from SIGMA (catalog number: B-5655).

Spots were counted on an Elispot reader (Bioreader 2000 from Biosys, Germany) in the size range between a diameter of 130 µm to 2000 µm. The average number of spontaneously induced IFN-γ spots was deduced from negative control samples (HCV or HIV peptide). Every response exceeding the average spontaneous IFN-γ release, to which value 2x the standard deviation of the negative control samples was added, was considered significant.

Collection, preparation and cryopreservation of PBMC for screening T cells responses against the CMV pp65 antigen
PBMC of 9 CMV seropositive HLA A2 healthy volunteers were included in this screen. HLA information of these donors on HLA class I A and B was available, but not on HLA class I C nor on HLA class II.

Whole blood was collected in ACD Vacutainer tubes (Becton Dickinson, Europe). PBMC were separated from whole blood on Lymphoprep (from Nycomed Pharma AS, Oslo, Norway) using Leuco Sep tubes (from Greiner), washed 3x with PBS (from GIBCOBRL, catalog number 14190-094) and resuspended at a concentration of 20 Mio c/ml in the following freezing medium: 4 parts RPMI 1640 from GIBCOBRL (catalog number: 31870-025) supplemented with 1 mM sodium pyruvate from GIBCOBRL (catalog number: 11360-039), 2 mM L-glutamine from GIBCOBRL (catalog number: 25030-024), 0,1 mM non-essential amino acids from GIBCOBRL (catalog number: 11140-035), 50 µg/ml gentamycin from GIBCOBRL (catalog number: 15710-049), 50 µM 2-mercaptoethanol from GIBCOBRL (catalog number: 31350-010) ; 5 parts fetal calf serum (FCS) from PAA (catalog number: A11-042); prior to use 1 part DMSO from SIGMA cell culture (catalog number D2650) was added.

PBMC were stored over night in freezing containers (Nalgene 1°C freezing container, catalog number 5100-0001) at -80°C and then transferred into the liquid nitrogen container.

- 35 -

ELISPOT Assay for single cell human IFN-gamma release: Detection of pp65-specific effectors from frozen PBMC

The assay was essentially done as described in Lalvani et al. Briefly, Multi Screen 96well filtration plates from Millipore (catalog-number:MAHA S4510) were coated with 10 µg/ml anti human IFN-γ mab B140 from Bender Med Systems (1 µg/well) over night at 4°C. Plates were washed 2 times with PBS (from GIBCOBRL, catalognumber 14190-094) and blocked with the following ELISPOT medium: RPMI 1640 from GIBCOBRL (catalog number: 31870-025) supplemented with 1 mM sodium pyruvate from GIBCOBRL (catalog number: 11360-039), 2 mM L-glutamine from GIBCOBRL (catalog number: 25030-024), 0,1 mM non-essential amino acids from GIBCOBRL (catalog number: 11140-035), 50 µg/ml gentamycin from GIBCOBRL (catalog number: 15710-049), 50 µM 2-mercaptoethanol from GIBCOBRL (catalog number: 31350-010) and 10% human serum type AB from BioWhittaker (catalog number: 14-490E) .

Cryopreserved PBMC were thawed quickly in a 37°C water bath and washed 2 with ELISPOT medium. Following thawing, PBMC were placed for 2h into an incubator (37°C, 5% CO₂). After this incubation, cells were filtered through a 70 µm cell strainer (Falcon), adjusted to a concentration of 2 Mio/ml and plated at 200.000 PBMC/well.

PBMC were cocultivated with either peptide pools, where each single peptide was contained at a final concentration of 5 µg/ml, or individual peptides at a final concentration of 10 µg/ml for 20 hrs. As an assay control polyclonal induction of IFN-γ secretion by Concanavalin A (SIGMA) was used. Spontaneous IFN-γ release was measured by either incubating PBMC with medium alone or -since exclusively HIV-negative donors were screened- by addition of an HLA 0201 restricted CTL epitope from HIV (HIV Reverse Transcriptase, aa 476-484: ILKEPVHGV). As positive control peptides frequently recognized HLA A 0201 restricted CTL epitopes from Influenza Matrixprotein (aa58-65: GILGFVFTL) and EBV BMLF1 antigen (aa 280-288: GLCTLVAML) were included in the screen.

Assays were developed as described in the paragraph "Synthetic

- 36 -

Elispot assays".

Immunization of HLA-transgenic mice

In the first set of experiments for immunogenicity the longer synthetic peptides incorporating candidate epitope sequences were injected into HLA-DRB1*0401-transgenic mice as follows: groups of 3 mice (female mice, 8 weeks of age) were injected subcutaneous into the hind footpad (in total 300µg of peptide and 5 nanomole of CpG1668 per mouse) once.

In the second set of experiments for epitope mapping synthetic CMV-derived peptides were tested in HLA-DRB1*0401-transgenic mice as follows: groups of 6 mice (female mice, 8 weeks of age) were injected subcutaneous into the flank (in total 100µg of peptide and 50 µl of CFA once and twice with 50µl of IFA per mouse) 3 times in weekly intervals.

In the third set of experiments for epitope mapping synthetic CMV-derived peptides were tested in HLA-DRB1*0401-transgenic mice as follows: groups of 8 mice (female mice, 8 weeks of age) were injected subcutaneous into the hind footpad (in total 300µg of peptide and 5 nanomole of CpG1668 per mouse for peptide no. 1500 and the same amount of peptide no.1503 and no. 1504 with 50 µl of IFA per mouse) once.

Isolation of murine splenocytes and separation of CD4+ /CD8+ T cells

Spleens were removed on day 7 after last injection and pooled together for each group. To prepare the single cell suspension spleens were smashed in DMEM medium supplemented with 5% FCS 2 mmole L-glutamine, 50µg-ml gentamycine, 1% sodium pyruvate, 0.1% 2-mercaptoethanol and 1% nonessential amino acids (PAA Laboratories, Linz, Austria) (complete medium), and filtered through a 70µm cell strainer. Cells were washed in complete medium (1200rpm, 10 minutes), the pellet was resuspended in red blood cell lysing buffer (Sigma-Aldrich) and incubated for 2 minutes to remove erythrocytes. After washing, cells were counted using KOVA Glasstic slides (Hycor, Biomedical INC.) and adjusted to

- 37 -

the concentration 1×10^7 ; 3.3×10^6 and 1.1×10^6 cells per ml with complete medium.

To determine class I or class II restriction of potential epitopes, total cells from murine spleens were separated into CD4+ and CD8+ populations by using MACS separation system with miniMACS and midiMACS columns and anti-CD4 and anti-CD8 magnetic beads (Miltenyi Biotec, Germany), according to supplier recommended procedure. The purity of CD4+ and CD8+ populations was checked by FACS analysis after staining shortly described as following: aliquots of cells 2×10^5 from total spleen cells, CD4+ and CD8+ fractions were stained with anti CD8-PE (53-6-7) and anti CD4-FITC (RM4.4) antibodies (PharMingen) for 15 minutes at room temperature. After washing, samples were analyzed on FACS CALIBUR (Becton Dickinson). Spleen cells from naïve DRB1*0401-transgenic mouse were stained in a similar way as experimental samples and additionally with isotype control antibodies, single and double stained samples to adjust FACS CALIBUR settings.

ELISPOT assay for IFN-gamma release from murine splenocytes

ELISpot assay plates (MAHA S4510, Millipore, Germany) were rinsed with PBS (200µl-well), coated with anti-mouse IFN-gamma (INF-γ) mAb (clone R46A2 purchased from ATCC, Manassas, VA; 50µl-well of 1µg/ml in 0.1 M NaHCO₃, pH 9.2-9.5) and incubated overnight at 4°C. Plates were washed four times with PBS containing 0.1% Tween-20 and incubated with PBS supplemented with 1%BSA (200µl/well) at room temperature for two hours to block nonspecific binding. Cells were seeded at total amount of 1×10^6 and 3.3×10^5 and 1.1×10^5 cells per well in 100µl and incubated overnight at 37°C/5%CO₂ with 10 µg/ml (final concentration) of different stimulants added individually to the wells containing cells in volume of 100µl: either long peptide used for vaccinations(relevant peptides) or overlapping 15-mers, derived from the longer peptide, or irrelevant peptide HA306-318 (no. 1171), not used for vaccination, or medium control. Subsequently, plates were washed four times and incubated with biotinylated anti-mouse IFN-γ mAb (clone AN18.17.24 purchased from ATCC, Manassas, VA; 100µl/well of 2µg/ml in PBS/1%BSA) for two hours at 37°C. After washing, 100µl/well of streptavidine-peroxidase di-

- 38 -

luted 1:5000 in PBS (Roche Diagnostics, Vienna, Austria) was added and plates were incubated at 37°C for one additional hour. Afterwards, plates were washed four times with PBS/0.1% Tween-20 and 100µl/well of the substrate (10ml of 10mM Tris pH 7.5 supplemented with 200µl of 40mg/ml DAB, 50ml of 80mg/ml NiCl₂ and 5 µl of 30% H₂O₂) was added. The reaction was stopped after 20-30 minutes by washing the plates with tap water. Dried plates were analyzed on BIOREADER 2000 (BioSys, Karben, Germany) and MICROSOFT OFFICE EXCEL program.

Example I. Epitope capture from peptide mixtures and identification by mass spectrometry

In this example the ability to capture high affinity peptides binding to HLA class II molecules from relatively simple peptide mixtures is demonstrated.

First, binding affinities of some individual peptides to soluble DRB1*0401 molecules were detected in a direct binding assay (Fig. 1). Peptide affinity was defined as high or low in comparison with binding of well-known "strong" binders, YAR and HA306-318 (Valli et al. (1993) in SDS-stability assay (Table 1). The binding affinity of 1242 peptide was considered to be the highest, comparable with affinity of YAR peptide.

Second, the ability to capture peptides binding to HLA class II molecules from the mixture of two high affinity ligands was tested. The binding reaction contained 1 µM of soluble DRB1*0401 molecules and 5 µM of each YAR and 1242 peptide. After the HLA-peptide complexes were formed, they were separated from the excess of free peptides by gel filtration chromatography. Fractions containing MHC molecules were collected. Bound peptides were eluted from the complexes and analyzed by mass-spectrometry. As the read-out, both tested high affinity ligands were revealed in the complexes with MHC molecules (Fig. 2).

Third, it was investigated if the binding capacity of the high affinity ligand would be influenced by the excess of low affinity peptide. For this purpose, binding reaction of 1µM DRB1*0401 molecules with 1242 peptide added in the concentration from 5 to

- 39 -

50 μM in the presence of constant amount (100 μM) of low affinity peptide 1236 was performed, so that the excess of 1236 over the 1242 peptide was 2-20 fold. Peptide binding was assessed by SDS-stability assay and verified by MS analysis (Fig.3). This experiment shows that 20-fold molar excess of low affinity peptide does not impede the formation of stable MHC-complexes with high affinity ligand.

Finally, binding of the pool of 10 peptides of different binding affinities was tested (see Table 1). The final concentration of each peptide in binding reaction was 5 μM , except 1236 added in 55 μM , meaning that the concentration of each individual binding peptide was 20-fold lower than final concentration of all peptides. DRB1*0401 molecules were added to 1 μM . The formation of MHC-peptide complexes was evaluated as described above, using single peptides as controls (Fig. 4). Peptides captured by DR4 molecules were purified and sequenced by MS analysis. The result of this analysis yielded the two peptides with highest binding affinities: 1242 and YAR (see Table 1). The peptides with moderate and/or low affinities were not found in the complexes, probably due to the high concentration of strong competitors used in this assay. But when the concentration of binding peptides was reduced to 0.5 μM such that DR4 molecules were added in the excess over each peptide then the peptide of moderate affinity (HA306-318) was determined. These results demonstrated the possibility of capturing more than one high affinity peptides as well as peptides of moderate binding affinities from the peptide mixtures.

Table 1

Pep- tide ID	Sequence	DR4 binding*	Reference
YAR	YARFQSQTTLKQKT	+++	Valli et al., 1993
HA ₃₀₆₋₃₁₈	YPKYVKQNTLKLAT	++	Valli et al., 1993
1235	IDELKTNSSLLTSILTYHVV	-	
1236	TGSGAGIAQAAAGTV	+	

- 40 -

1237	GVSTANATVYMIDSVL	++	
1238	NFAGIEAAASAIQGNV	-	
1239	AETPGCVAYIGISFLDQ	-	
1240	VSDLKSSTAVIPGYPV	+	
1241	NFLLPDAQSIQAAAAG	++	
1242	YNINISLPSYYPDQKSL	+++	

"+++" high binding affinity; "++" moderate binding affinity; "+" low binding affinity; "-" no binding.

Example II. Measuring T cell responses against captured peptides by synthetic T cell assays

In this example it is demonstrated, that peptides identified as described in Example I, not only bind to MHC molecules, but are also capable of stimulating T cells, proofing, that they are T cell epitopes. This question is relevant, because binding of a peptide to a MHC molecule by itself does not guarantee, that this peptide is relevant "in vivo". The peptide might not be generated during "in vivo" antigen processing by the endosomal proteases (antigens for CD4+ T cells) or the proteasome (antigens for CD8+ T cells). And even peptides, which are processed and presented "in vivo" can instead of stimulating a T cell anergize it (antagonists). As a read out for T cell stimulation an IFN- γ ELISPOT assay was chosen.

For the example shown in Fig. 5a, the CD4+ T cells from healthy HCV-negative, BCG vaccinated donors were isolated from peripheral blood and cocultivated with SHLA DRB1* 0401 loaded with p1242 and p84, an HCV peptide, which was used as a negative control. Costimulation was provided by adding an anti CD28 monoclonal antibody. Fig. 5a shows the number of induced IFN- γ spots, each spot representing a stimulated T cell. The induced number is significantly above the spontaneous IFN- γ secretion detected in the sample with the negative control peptide.

In order to show, that this not only holds true for MHC class II restricted peptides, the CD8+ T cells from healthy, HIV negative EBV-infected donors with a recent Influenza infection were in-

- 41 -

cubated with sHLA A0201 molecules, loaded with either a peptide from the Influenza Matrix protein or the EBV BMLF1 antigen. Again a marked T cell response against both viral peptides could be detected (Fig 5b). The specificity of this response was proven by incubating the loaded sHLA A0201 molecules with anti HLA A0201 antibody prior to cocultivation with the CD8+ T cells. This preincubation nearly totally abolished IFN- γ secretion, only spontaneous IFN- γ secretion was detected in these samples. These examples demonstrate, that functional T cells against the captured peptides exist "in vivo", that these peptides are therefore epitopes and might be useful in a vaccine, inducing protective immune responses.

Example III. Rapid identification of HLA-binding peptides by measuring peptide pools arrayed in matrix format

After confirming the ability to use peptide pools for detection of individual binding peptides, the approach according to the present invention was applied to identify peptides capable to bind HLA class II molecules from CMV pp65 antigen. For this purpose, the direct peptide binding method was combined with the peptide pool array approach, described early (Kern et al., 1999; Tobery, et al., 2001).

In order to span the entire pp65 sequence of 560 aa long, 547 15 mers, overlapping by 14 aa, were designed. Sequences of all peptides are shown in Table 2.

Replacement Sheet

- 42 -

Table 2. Sequences of overlapping 15-mer peptides spanning the entire CMV pp65 sequence

Peptide no	Sequence	Peptide no	Sequence	Peptide no	Sequence
1	MESRRRCPEMISVL	83	TGSEVENSVNVHNHP	165	GLAWTRQQNQWKEPD
2	ESRRRCPEMISVLG	84	GSEVENSVNVHNPT	166	LAWTRQQNQWKEPDV
3	SRRRRCPEMISVLGP	85	SEVENSVNVHNPTG	167	AWTRQQNQWKEPDVY
4	RRRRCPEMISVLGPIS	86	EVENSVNVHNPTGR	168	WTRQQNQWKEPDVYY
5	GRRRCPEMISVLGPIS	87	VENSVNVHNPTGRS	169	TRQQNQWKEPDVYYT
6	RRCPEMISVLGPISGH	88	ENSVNVHNPTGRSIC	170	RQQNQWKEPDVYYTS
7	RCPEMISVLGPISGHV	89	NVSVNVHNPTGRSIC	171	QQNQWKEPDVYYTSA
8	CPEMISVLGPISGHVL	90	VSVNVHNPTGRSICP	172	QNQWKEPDVYYTSAF
9	PEMISVLGPISGHVLK	91	SVNVHNPTGRSICPS	173	NQWKEPDVYYTSAFV
10	EMISVLGPISGHVLK	92	VNVHNPTGRSICPSQ	174	QWKEPDVYYTSAFVF
11	MISVLGPISGHVLKAV	93	NVHNPTGRSICPSQE	175	WKEPDVYYTSAFVFP
12	ISVLGPISGHVLKAVF	94	VHNPTGRSICPSQEP	176	KEPDVYYTSAFVFPT
13	SVLGPISGHVLKAVFS	95	HNPTGRSICPSQEPMS	177	EPDVYYTSAFVFPTK
14	VLGPISGHVLKAVFSR	96	NPTGRSICPSQEPMSI	178	PDVYYTSAFVFPTKD
15	LGPISGHVLKAVFSRG	97	PTGRSICPSQEPMSIY	179	DVYYTSAFVFPTKDV
16	GPISGHVLKAVFSRGD	98	TGRSICPSQEPMSIYV	180	VYYTSAFVFPTKDVA
17	PISGHVLKAVFSRGDT	99	GRSICPSQEPMSIYVY	181	YYTSAFVFPTKDVAL
18	ISGHVLKAVFSRGDTP	100	RSICPSQEPMSIYVYA	182	YTSAFVFPTKDVALR
19	SGHVLKAVFSRGDTPV	101	SICPSQEPMSIYVYAL	183	TSAFVFPTKDVALRH
20	GHVLKAVFSRGDTPVL	102	ICPSQEPMSIYVYALP	184	SAFVFPTKDVALRHV
21	HVLKAVFSRGDTPVLP	103	CPSQEPMSIYVYALPL	185	AFVFPTKDVALRHVV
22	VLKAVFSRGDTPVLPHE	104	PSQEPMSIYVYALPLK	186	FVFPTKDVALRHHVC
23	LKAVFSRGDTPVLPHE	105	QEPMSIYVYALPLKML	187	VFPTKDVALRHHVCA
24	KAVFSRGDTPVLPHE	106	QEPMSIYVYALPLKMLN	188	FPTKDVALRHHVCAH
25	AVFSRGDTPVLPHE	107	EPMSIYVYALPLKMLNI	189	PTKDVALRHHVCAHE
26	VFSRGDTPVLPHE	108	PMSIYVYALPLKMLNIP	190	TKDVALRHHVCAHEL
27	FSRGDTPVLPHE	109	SIYVYALPLKMLNIPSI	191	KDVALRHHVCAHELVC
28	SRGDTPVLPHE	110	IYVYALPLKMLNIPSI	192	DVALRHHVCAHELVC
29	RGDTPVLPHE	111	YVYALPLKMLNIPSI	193	VALRHHVCAHELVC
30	GDTPLVLPHE	112	VYALPLKMLNIPSI	194	ALRHHVCAHELVC
31	DTPVLPHE	113	YALPLKMLNIPSI	195	LRHHVCAHELVC
32	TPVLPHE	114	ALPLKMLNIPSI	196	RHHVCAHELVC
33	PVLPHE	115	ALPLKMLNIPSI	197	HVCAHELVC
34	VLPHETRL	116	LPLKMLNIPSI	198	VVCAHELVC
35	LPHETRL	117	PLKMLNIPSI	199	VCAHELVC
36	PHETRL	118	LKMLNIPSI	200	CAHELVC
37	HETRL	119	KMLNIPSI	201	AHELVC
38	ETRL	120	MLNIPSI	202	HELVC
39	TRL	121	LNIPSI	203	ELVC
40	RL	122	NIPSI	204	LVC
41	LL	123	IPSI	205	VCS
42	L	124	PSI	206	CS
43	Q	125	SIN	207	SM
44	T	126	INV	208	MENT
45	G	127	NV	209	ENT
46	I	128	VH	210	N
47	H	129	HH	211	TR
48	V	130	HY	212	R
49	R	131	PS	213	AT
50	V	132	SA	214	TK
51	S	133	AA	215	K
52	Q	134	AE	216	M
53	P	135	ER	217	Q
54	S	136	K	218	V
55	L	137	R	219	I
56	I	138	K	220	G
57	L	139	H	221	D
58	V	140	R	222	Q

Replacement Sheet

- 43 -

59	SOYTPDSTPCHRGDN	141	HLPVADAVIHASGKQ	223	YVKVYLESFCEDVPS
60	QYTPDSTPCHRGDNQ	142	LPVADAVIHASGQM	224	VKVYLESFCEDVPSG
61	YTPDSTPCHRGDNQL	143	PVADAVIHASGQMWW	225	KVYLESFCEDVPSGK
62	TPDSTPCHRGDNQLQ	144	VADAVIHASGQMWWQ	226	VYLESFCEDVPSGKL
63	PDSTPCHRGDNQLQV	145	ADAVIHASGQMWWQA	227	YLESFCEDVPSGKLF
64	DSTPCHRGDNQLQVQ	146	DAVIHASGQMWWQAR	228	LESFCEDVPSGKLFM
65	STPCHRGDNQLQVQH	147	AVIHASGQMWWQARL	229	ESFCEDVPSGKLFMH
66	TPCHRGDNQLQVQHT	148	VIHASGQMWWQARLT	230	SFCEDVPSGKLFMHV
67	PCHRGDNQLQVQHTY	149	IHASGQMWWQARLTV	231	FCEDVPSGKLFMHVT
68	CHRGDNQLQVQHTYF	150	HASGQMWWQARLTVS	232	CEDVPSGKLFMHVTL
69	HRGDNQLQVQHTYFT	151	ASGQMWWQARLTVSG	233	EDVPSGKLFMHVTLG
70	RGDNQLQVQHTYFTG	152	SGQMWWQARLTVSGL	234	DVPSGKLFMHVTLGS
71	GDNLQVQHTYFTGSE	153	GQMWWQARLTVSGLA	235	VPSGKLFMHVTLGSD
72	DNQLQVQHTYFTGSE	154	QMWQARLTVSGLAW	236	PSGKLFMHVTLGSDV
73	NQLQVQHTYFTGSEV	155	QMWQARLTVSGLAWT	237	SGKLFMHVTLGSDVE
74	QLQVQHTYFTGSEVE	156	MWQARLTVSGLAWTR	238	GKLFMHVTLGSDVEE
75	LQVQHTYFTGSEVEN	157	WQARLTVSGLAWTRQ	239	KLFMHVTLGSDVEED
76	QVQHTYFTGSEVENV	158	QARLTVSGLAWTRQQ	240	LFMHVTLGSDVEEDL
77	VQHTYFTGSEVENVS	159	ARLTVSGLAWTRQQN	241	FMHVTLGSDVEEDLT
78	QHTYFTGSEVENVSV	160	RLTVSGLAWTRQQNQ	242	MHVTLGSDVEEDLTM
79	HTYFTGSEVENVSVN	161	LTVSGLAWTRQQNQW	243	HVTLGSDVEEDLTM
80	TYFTGSEVENVSVNV	162	TVSGLAWTRQQNQWK	244	VTGSDVEEDLTMTR
81	YFTGSEVENVSVNVH	163	VSGLAWTRQQNQWKE	245	TLGSDVEEDLTMTRN
82	FTGSEVENVSVNVHN	164	SGLAWTRQQNQWKEP	246	LGSDVEEDLTMTRNP

Peptide no	Sequence	Peptide no	Sequence	Peptide no	Sequence
247	GSDVEEDLTMTRNPQ	329	EVQAIRETVELRQYD	411	VTTERKTTPRVTTGGGA
248	SDVEEDLTMTRNPQP	330	VQAIRETVELRQYDP	412	TTERKTTPRVTTGGGAM
249	DVEEDLTMTRNPQPF	331	QAIRETVELRQYDPV	413	TERKTTPRVTTGGGAMA
250	VEEDLTMTRNPQPFFM	332	AIRETVELRQYDPVA	414	ERKTTPRVTTGGGAMAG
251	EEDLTMTRNPQPFFMR	333	IRETVELRQYDPVAA	415	RKTTPRVTTGGGAMAGA
252	EDLTMTRNPQPFFMRP	334	RETVELRQYDPVAAL	416	KTPRVTTGGGAMAGAS
253	DLTMTRNPQPFFMRPH	335	ETVELRQYDPVAALF	417	TPRVTTGGGAMAGAST
254	LTMTRNPQPFFMRPHE	336	TVELRQYDPVAALFF	418	PRVTGGGAMAGASTS
255	TMTRNPQPFFMRPHER	337	VELRQYDPVAALFFF	419	RVTGGGAMAGASTSA
256	MTRNPQPFFMRPHERN	338	ELRQYDPVAALFFFD	420	VTGGGAMAGASTSAG
257	TRNPQPFFMRPHERNG	339	LRQYDPVAALFFFDI	421	TGGGAMAGASTSAGR
258	RNPQPFFMRPHERNGF	340	RQYDPVAALFFFDID	422	GGGAMAGASTSAGRK
259	NPQPFFMRPHERNGFT	341	QYDPVAALFFFDIDL	423	GGAMAGASTSAGRKR
260	PQPFFMRPHERNGFTV	342	YDPVAALFFFDIDLL	424	GAMAGASTSAGRKRK
261	QPFFMRPHERNGFTVL	343	DPVAALFFFDIDLLL	425	AMAGASTSAGRKRKS
262	PFFMRPHERNGFTVLC	344	PVAALFFFDIDLLLQ	426	MAGASTSAGRKRKSA
263	FMRPHERNGFTVLC	345	VAALFFFDIDLLLQR	427	AGASTSAGRKRKSAS
264	MRPHERNGFTVLC	346	AALFFFDIDLLLQRG	428	GASTSAGRKRKSASS
265	RPHERNFTVLC	347	ALFFFDIDLLLQRGP	429	ASTSAGRKRKSASSA
266	PHERNFTVLC	348	LFFFDIDLLLQRGPQ	430	STSAGRKRKSASSAT
267	HERNGFTVLC	349	FFFDIDLLLQRGPQY	431	TSAGRKRKSASSATA
268	ERNNGFTVLC	350	FFDIDLLLQRGPQYS	432	SAGRKRKSASSATAC
269	RNGFTVLC	351	FDIDLLLQRGPQYSE	433	AGRKRKSASSATACT
270	NGFTVLC	352	DIDLLLQRGPQYSEH	434	GRKRKSASSATACT
271	GFTVLC	353	IDLLLQRGPQYSEHP	435	RKRKSASSATACTSG
272	FTVLC	354	DLLLQRGPQYSEHPT	436	KRKSASSATACTSGV
273	TVLC	355	LLLQRGPQYSEHPTF	437	RKSASSATACTSGVM
274	VLC	356	LLQRGPQYSEHPTFT	438	KSASSATACTSGVMT
275	LCPKNMIKPGKISH	357	LQRGPQYSEHPTFTS	439	SASSATACTSGVMTR
276	CPKNMIKPGKISHI	358	QRGPQYSEHPTFTSQ	440	ASSATACTSGVMTRG
277	PKNMIKPGKISHIM	359	RGPQYSEHPTFTSQY	441	SSATACTSGVMTRGR
278	KNMIKPGKISHIML	360	GPQYSEHPTFTSQYR	442	SATACTSGVMTRGRRL
279	NMIKPGKISHIMLD	361	PQYSEHPTFTSQYRI	443	ATACTSGVMTRGRRLK
280	MIKPGKISHIMLDV	362	QYSEHPTFTSQYRIQ	444	TACTSGVMTRGRRLKA
281	IIPKPGKISHIMLDVA	363	YSEHPTFTSQYRIQG	445	ACTSGVMTRGRRLKAE

Replacement Sheet

- 44 -

282	IKPGKISHIMLDVAF	364	SEHPTFTSQYRIQGK	446	CTSGVMTRGRLKAES
283	KPGKISHIMLDVAF	365	EHPTFTSQYRIQGKL	447	TSGVMTRGRLKAEST
284	PGKISHIMLDVAF	366	HPTFTSQYRIQGKLE	448	SGVMTRGRLKAESTV
285	GKISHIMLDVAF	367	PTFTSQYRIQGKLEY	449	GVMTRGRLKAESTVA
286	KISHIMLDVAF	368	TFTSQYRIQGKLEYR	450	VMTRGRLKAESTVAP
287	ISHIMLDVAF	369	FTSQYRIQGKLEYRH	451	MTRGRLKAESTVAPE
288	SHIMLDVAF	370	TSQYRIQGKLEYRHT	452	TRGRLKAESTVAPEE
289	HIMLDVAF	371	SQYRIQGKLEYRHTW	453	RGRKAESTVAPEED
290	IMLDVAF	372	QYRIQGKLEYRHTWD	454	GRKAESTVAPEEDT
291	MLDVAF	373	YRIQGKLEYRHTWDR	455	RLKAESTVAPEEDTD
292	LDVAF	374	RIQGKLEYRHTWDRH	456	LKAESTVAPEEDTDE
293	DVAF	375	IQGKLEYRHTWDRHD	457	KAESTVAPEEDTDED
294	VAF	376	QGKLEYRHTWDRHDE	458	AESTVAPEEDTDEDS
295	AFT	377	GKLEYRHTWDRHDEG	459	ESTVAPEEDTDESD
296	FTS	378	KLEYRHTWDRHDEGA	460	STVAPEEDTDESDN
297	TS	379	LEYRHTWDRHDEGAA	461	TVAPEEDTDESDNE
298	S	380	EYRHTWDRHDEGAAQ	462	VAPEEDTDESDNEI
299	HE	381	YRHTWDRHDEGAAQG	463	APPEEDTDESDNEIH
300	EH	382	RHTWDRHDEGAAQGD	464	PEEDTDESDNEIHN
301	HF	383	HTWDRHDEGAAQGDD	465	EEDTDESDNEIHN
302	FG	384	TWDRHDEGAAQGDDDD	466	EDTDESDNEIHNPA
303	G	385	WDRHDEGAAQGDDDV	467	DTDESDNEIHNPAV
304	LL	386	DRHDEGAAQGDDDVW	468	TDESDNEIHNPAVF
305	LC	387	RHDEGAAQGDDDVWT	469	DESDNEIHNPAVFT
306	CP	388	HDEGAAQGDDDVWTS	470	EDSDNEIHNPAVFTW
307	PK	389	DEGAAQGDDDVWTSG	471	DSDNEIHNPAVFTWP
308	KS	390	EGAAQGDDDVWTSGS	472	SDNEIHNPAVFTWPP
309	SIP	391	GAAQGDDDVWTSGSD	473	DNEIHNPAVFTWPPW
310	IP	392	AAQGDDDVWTSGSDS	474	NEIHNPAVFTWPPWQ
311	P	393	AQGDDDVWTSGSDSD	475	EIHNPAVFTWPPWQA
312	GL	394	QGDDDVWTSGSDSDE	476	IHNPAVFTWPPWQAG
313	LS	395	GDDDVWTSGSDSDEE	477	HNPVFTWPPWQAGI
314	SI	396	DDDVWTSGSDSDEEL	478	NPAVFTWPPWQAGIL
315	IS	397	DDVWTSGSDSDEELV	479	PAVFTWPPWQAGILA
316	SG	398	DVWTSGSDSDEELVT	480	AVFTWPPWQAGILAR
317	GN	399	VWTSGSDSDEELVTT	481	VFTWPPWQAGILARN
318	N	400	WTSGSDSDEELVTTE	482	FTWPPWQAGILARNL
319	LL	401	TSGSDSDEELVTTER	483	TWPPWQAGILARNLV
320	LM	402	SGSDSDEELVTTERK	484	WPPWQAGILARNLVP
321	MN	403	GSDSDEELVTTERKT	485	PPWQAGILARNLVPM
322	NG	404	SDSDEELVTTERKTP	486	PWQAGILARNLVPMV
323	GQ	405	DSDEELVTTERKTPR	487	WQAGILARNLVPMVA
324	QQ	406	SDEELVTTERKTPRV	488	QAGILARNLVPMVAT
325	QI	407	DEELVTTERKTPRVT	489	AGILARNLVPMVATV
326	IF	408	EELVTTERKTPRVTG	490	GILARNLVPMVATVQ
327	F	409	ELVTTERKTPRVTGG	491	ILARNLVPMVATVQG
328	LEV	410	LVTTERKTPRVTGGG	492	LARNLVPMVATVQGG

Peptide no Sequence
 493 ARNLVPMVATVQGGN
 494 RNLVPMVATVQGGNL
 495 NLVPMVATVQGGNLK
 496 LVPMVATVQGGNLKY
 497 VPMVATVQGGNLKYQ
 498 PMVATVQGGNLKYQE
 499 MVATVQGGNLKYQEF
 500 VATVQGGNLKYQEFF
 501 ATVQGGNLKYQEFFW
 502 TVQGGNLKYQEFFWD
 503 VQGGNLKYQEFFWDA

Peptide no Sequence
 520 TYRIFAELEGVWQPA
 521 YRIFAELEGVWQPAA
 522 RIFAELEGVWQPAAQ
 523 IFAELEGVWQPAAQP
 524 FAELEGVWQPAAQPK
 525 AELEGVWQPAAQPKR
 526 ELEGVWQPAAQPKRR
 527 LEGVWQPAAQPKRRR
 528 EGVWQPAAQPKRRRH
 529 GVWQPAAQPKRRRHR
 530 VWQPAAQPKRRRHRQ

Peptide no Sequence
 544 QDALPGPCIASTPKK
 545 DALPGPCIASTPKKH
 546 ALPGPCIASTPKKHR
 547 LPGPCIASTPKKHRG

Replacement Sheet

- 44a -

504	QGQNLKYQEFFWDAN	531	WQPAAQPKRRRHRQD
505	GQNLKYQEFFWDAND	532	QPAAQPKRRRHRQDA
506	QNLKYQEFFWDANDI	533	PAAQPKRRRHRQDAL
507	NLKYQEFFWDANDIY	534	AAQPKRRRHRQDALP
508	LKYQEFFWDANDIYR	535	AQPKRRRHRQDALPG
509	KYQEFFWDANDIYRI	536	QPKRRRHRQDALPGP
510	YQEFFWDANDIYRIF	537	PKRRRHRQDALPGPC
511	QEFFWDANDIYRIFA	538	KRRRHRQDALPGPCI
512	EFFWDANDIYRIFAE	539	RRRHRQDALPGPCIA
513	FFWDANDIYRIFAE	540	RRHRQDALPGPCIAS
514	FWDANDIYRIFAELE	541	RHRQDALPGPCIAST
515	WDANDIYRIFAELEG	542	HRQDALPGPCIASTP
516	DANDIYRIFAELEGV	543	RQDALPGPCIASTPK
517	ANDIYRIFAELEGVW	544	QDALPGPCIASTPKK
518	NDIYRIFAELEGVWQ	545	DALPGPCIASTPKKH
519	DIYRIFAELEGVWQP	546	ALPGPCIASTPKKHR
520	IYRIFAELEGVWQPA	547	LPGPCIASTPKKHRG
521	YRIFAELEGVWQPAA	519	DIYRIFAELEGVWQP
522	RIFAELEGVWQPAAQ	520	IYRIFAELEGVWQPA
523	IFAELEGVWQPAAQP	521	YRIFAELEGVWQPAA
524	FAELEGVWQPAAQPK	522	RIFAELEGVWQPAAQ
525	AELEGVWQPAAQPKR	523	IFAELEGVWQPAAQP
526	ELEGVWQPAAQPKRR	524	FAELEGVWQPAAQPK
527	LEGVWQPAAQPKRRR	525	AELEGVWQPAAQPKR
528	EGVWQPAAQPKRRRH	526	ELEGVWQPAAQPKRR
529	GVWQPAAQPKRRRHR	527	LEGVWQPAAQPKRRR
530	VWQPAAQPKRRRHRQ	528	EGVWQPAAQPKRRRH
531	WQPAAQPKRRRHRQD	529	GVWQPAAQPKRRRHR
532	QPAAQPKRRRHRQDA	530	VWQPAAQPKRRRHRQ
533	PAAQPKRRRHRQDAL	531	WQPAAQPKRRRHRQD
534	AAQPKRRRHRQDALP	532	QPAAQPKRRRHRQDA
535	AQPKRRRHRQDALPG	533	PAAQPKRRRHRQDAL
536	QPKRRRHRQDALPGP	534	AAQPKRRRHRQDALP
537	PKRRRHRQDALPGPC	535	AQPKRRRHRQDALPG
538	KRRRHRQDALPGPCI	536	QPKRRRHRQDALPGP
539	RRRHRQDALPGPCIA	537	PKRRRHRQDALPGPC
540	RRHRQDALPGPCIAS	538	KRRRHRQDALPGPCI
541	RHRQDALPGPCIAST	539	RRRHRQDALPGPCIA
542	HRQDALPGPCIASTP	540	RRHRQDALPGPCIAS
543	RQDALPGPCIASTPK	541	RHRQDALPGPCIAST
544	QDALPGPCIASTPKK	542	HRQDALPGPCIASTP
545	DALPGPCIASTPKKH	543	RQDALPGPCIASTPK
546	ALPGPCIASTPKKHR	544	QDALPGPCIASTPKK
547	LPGPCIASTPKKHRG	545	DALPGPCIASTPKKH
493	ARNLVPVATVQGQN	546	ALPGPCIASTPKKHR
494	RNLVPMVATVQGQNL	547	LPGPCIASTPKKHRG
495	NLVPVATVQGQNLK	519	DIYRIFAELEGVWQP
496	LVPVATVQGQNLKY	520	IYRIFAELEGVWQPA
497	VPMVATVQGQNLKYQ	521	YRIFAELEGVWQPAA
498	PMVATVQGQNLKYQE	522	RIFAELEGVWQPAAQ
499	MVATVQGQNLKYQEF	523	IFAELEGVWQPAAQP
500	VATVQGQNLKYQEFF	524	FAELEGVWQPAAQPK
501	ATVQGQNLKYQEFFW	525	AELEGVWQPAAQPKR
502	TVQGQNLKYQEFFWD	526	ELEGVWQPAAQPKRR
503	VQGQNLKYQEFFWDA	527	LEGVWQPAAQPKRRR
504	QGQNLKYQEFFWDAN	528	EGVWQPAAQPKRRRH
505	GQNLKYQEFFWDAND	529	GVWQPAAQPKRRRHR
506	QNLKYQEFFWDANDI	530	VWQPAAQPKRRRHRQ
507	NLKYQEFFWDANDIY	531	WQPAAQPKRRRHRQD
508	LKYQEFFWDANDIYR	532	QPAAQPKRRRHRQDA
509	KYQEFFWDANDIYRI	533	PAAQPKRRRHRQDAL
510	YQEFFWDANDIYRIF	534	AAQPKRRRHRQDALP

Replacement Sheet

- 44b -

511	QEFFWDANDIYRIFA	535	AQPKRRRHRQDALPG
512	EFFWDANDIYRIFAE	536	QPKRRRHRQDALEGP
513	FFWDANDIYRIFAE	537	PKRRRHRQDALPGPC
514	FWDANDIYRIFAELE	538	KRRRHRQDALPGPCI
515	WDANDIYRIFAELEG	539	RRRHRQDALPGPCIA
516	DANDIYRIFAELEGV	540	RRHRQDALPGPCIAS
517	ANDIYRIFAELEGVW	541	RHRQDALPGPCTAST
518	NDIYRIFAELEGVWQ	542	HRQDALPGPCTASTP
519	DIYRIFAELEGVWQP	543	RQDALPGPCTASTPK

- 45 -

Because of the huge peptide number to be tested, peptides were combined in 42 pools each containing 21 peptides arrayed in a matrix format, as shown on Fig. 6. The mixtures were prepared such that each individual peptide was contained in exactly one "row" pool and one "column" pool.

Each matrix pool was tested for binding affinity to DR4 molecules in SDS-stability assay; HA306-318 was used as a reference peptide (Fig. 7a). In the first screen 18 out of 42 positive peptide pools were found : no. 2,3,6,15,16,19,20 from "row" pools and no. 28-38 from "column" pools (see Fig. 6). By finding the intersections of reactive pools in the array, the 77 peptides were determined as potential binders. Then each single 15 mer was checked for binding to DR4 molecules individually (Fig. 7b, Table 3). As a result, 20 peptides selected in the first screen were confirmed to be binders in the second screen. Usually, in rows of the array more than one positive peptides was found. Probably, these peptides overlapping by 14 aa represent longer epitopes recognized by TCR receptors on CD4+ T cells.

Four peptides (peptide no. 177, 361,507,508) containing partially or fully identical sequences were described earlier as capable to evoke specific CD4+ T cells response in healthy, CMV-exposed human subjects (Khattab et al., 1997; Bitmansour et al., 2001). This confirms the validity of our approach.

Some of the identified DRB1*0401 binding peptides, as well as others chosen by prediction algorithms were also tested for binding to soluble HLA molecules of different alleles: DRB1*0101, DRB1*0404, DRB1*0701 and DRB1*1101 (Table 3). Seven peptides (No 58, 177, 559, 360, 452, 470 and No 496) were demonstrated to have affinity for at least two HLA molecules arguing for promiscuity of these epitopes.

Three peptides binding to DRB1*0404 (no. 421, 469 and 470) were also found to induce IFN-gamma secretion in T cells from CMV seropositive donors (see Example IV) again indicating that also this second reverse immunological approach can identify true T cell epitopes.

Replacement Sheet

- 46 -

In conclusion, several novel sequences derived from the CMV pp65 antigen, competent of binding to HLA class II molecules, considered of being candidates for class II epitopes (shaded in Table 3) were identified. Immunogenicity of peptides, incorporating mentioned epitopes, as well as their CD4+ specificity was verified by testing their ability to induce IFN-gamma production upon ex vivo stimulation of T cell from CMV seropositive donors and/or HLA-DRB1*0401 transgenic mice (see Example IV, VI and VII).

Table 3.

Binding of individual 15mer peptides of CMV pp65 protein to soluble HLA-molecules. Newly identified binders are shaded.

Mixture No.	Peptide No	Peptide sequence	Binding# to DRB1 *0401	Binding to DRB1 *0404	Binding to DRB1 *0101	Reference, comment
2 and 31	37	HETRLQLQTGIHVRVS	-	nd	nd	Gallot et al., CD4+ clone / DQ*0602
2 and 35	41	LLQTGIHVRVSQPSL	-	nd	nd	Gallot et al., CD4+ clone / DQ*0602
2 and 37	43	QTGIHVRVSQPSLIL	-	nd	nd	Gallot et al., CD4+ clone / DQ*0602
3 and 27	54	SLILVSQYTPDSTPC	-	nd	nd	
3 and 28	55	LILVSQYTPDSTPCH	+	nd	nd	new, claim
3 and 29	56	ILVSQYTPDSTPCHR	++	nd	nd	new, claim
3 and 30	57	LVSQYTPDSTPCHRG	++	-	-	new, claim
3 and 31	58	VSQYTPDSTPCHRGD	++	nd	nd	new, claim
3 and 32	59	SQYTPDSTPCHRGDN	++	nd	nd	new, claim
3 and 33	60	QYTPDSTPCHRGDNQ	++	nd	nd	new, claim

Replacement Sheet

- 47 -

Mixture No.	Peptide No	Peptide sequence	Binding# to DRB1 *0401	Binding to DRB1 *0404	Binding to DRB1 *0101	Reference, comment
3 and 34	61	YTPDSTPCHRGDNQL	++	nd	nd	new, claim
3 and 35	62 ^s	TPDSTPCHRGDNQLQ	-	nd	nd	new, claim
3 and 36	63 ^s	PDSTPCHRGDNQLQV	-	nd	nd	new, claim
3 and 37	64	DSTPCHRGDNQLQVQ	+	nd	nd	new, claim
3 and 38	65	STPCHRGDNQLQVQH	-	nd	nd	
4 and 23	107	EPMSIYVYALPLKML	-	nd	-	
4 and 25	109	SIYVYALPLKMLNIP	-	nd	+	new, claim
4 and 27	111	IYVYALPLKMLNIPS	-	nd	-	
6 and 29	170	RQQNQWKEPDVYYTS	-	nd	nd	
6 and 31	172	QNQWKEPDVYYTSAF	-	nd	nd	
6 and 33	174	QWKEPDVYYTSAFVF	-	nd	-	
6 and 34	175	WKEPDVYYTSAFVFP	-	nd	+	
6 and 35	176	KEPDVYYTSAFVFPT	+/-	nd	-	
6 and 36	177	EPDVYYTSAFVFPTK	+	nd	nd	Bitmansour et al., CD4+ clone
6 and 37	178	PDVYYTSAFVFPTKD	++	nd	nd	
6 and 38	179	DVYYTSAFVFPTKDV	+	nd	nd	
6 and 39	180	VYYTSAFVFPTKDVA	+/-	nd	nd	
15 and 27	357	LQRGPOYSEHPTFTTS	-	nd	nd	
15 and 28	358	QRGPOYSEHPTFTSQ	+	nd	nd	
15 and 29	359	RGPOYSEHPTFTSQY	+/-	nd	nd	
15 and 30	360	GPQYSEHPTFTSQYR	+/-	nd	nd	
15 and 31	361	PQYSEHPTFTSQYRI	+/-	nd	nd	Khattab et al., CD4+ clone / DR11
15 and 32	362	QYSEHPTFTSQYRIQ	-	nd	nd	

Replacement Sheet

- 48 -

Mixture No.	Peptide No	Peptide sequence	Binding# to DRB1 *0401	Binding to DRB1 *0404	Binding to DRB1 *0101	Reference, comment
15 and 33	363	YSEHPTFTSQYRIQG	-	nd	nd	
15 and 35	365	EHPTFTSQYRIQGKL	-	nd	nd	Gallot et al., CD4+ clone / DR*1302
15 and 37	367	PTFTSQYRIQGKLEY	-	nd	nd	Gallot et al., CD4+ clone / DR*1302
16 and 28	379	LEYRHTWDRHDEGAA	-	nd	nd	
16 and 29	380	EYRHTWDRHDEGAAQ	-	nd	nd	
16 and 31	382	RHTWDRHDEGAAQGD	-	nd	nd	
16 and 32	383	HTWDRHDEGAAQGDD	+/-	nd	nd	new, claim
16 and 33	384	TWDRHDEGAAQGDDD	+	nd	nd	new, claim
16 and 34	385	WDRHDEGAAQGDDDV	-	nd	nd	
16 and 37	388	HDEGAAQGDDDVWTS	-	nd	nd	
18 and 26	419	RVTGGGAMAGASTSA	-	-	nd	Bitmansour, CD4+ clone
18 and 28	421	TGGGAMAGASTSAGR	-	+	nd	new, claim
18 and 30	423	GGAMAGASTSAGRKR	-	-	nd	
nd	448	SGVMTRGRLKAESTV	-	nd	nd	
nd	449	GVMTRGRLKAESTVA	+/-	nd	nd	new, claim
nd	450	VMTRGRLKAESTVAP	+	nd	nd	new, claim

Replacement Sheet

- 49 -

Mixture No.	Peptide No	Peptide sequence	Binding# to DRB1 *0401	Binding to DRB1 *0404	Binding to DRB1 *0101	Reference, comment
nd	451	MTGRRLKAESTVAPE	+	nd	nd	new, claim
nd	452	TRGRRLKAESTVAPEE	+	+	nd	new, claim
nd	453	RGRLKAESTVAPEED	+	nd	nd	new, claim
nd	454	GRLKAESTVAPEEDT	+	nd	nd	new, claim
nd	455	RLKAESTVAPEEDTD	-	nd	nd	
18 and 35	468	TDSDNEIHNPAVF	-	-	nd	
18 and 36	469	DESDNEIHNPAVFT	-	+/-	nd	new, claim
18 and 37	470	EDSDNEIHNPAVFTW	-	+/-	nd	new, claim
19 and 38	492	LARNLVPMVATVQGQ	-	++	-	Bitmansour , CD4+ clone
19 and 40	494	RNLVPMVATVQGQNL	-	++	-	?
19 and 42	496	LVPVATVQGQNLKY	-	+	+	?
20 and 32	507	NLKYQEFFWDANDIY	+/-	nd	nd	Bitmansour et al. , CD4+ clone
20 and 33	508	LKYQEFFWDANDIYR	+/-	nd	nd	Bitmansour , Khattab, CD4+ / DR3
20 and 35	510	YQEFFWDANDIYRIF	-	nd	nd	Khattab et al., CD4+ / DR3
20 and 37	512	EFFWDANDIYRIFAE	-	nd	nd	

"-" no binding; "+/-" weak binding; "+" intermediate binding;

Replacement Sheet

- 50 -

"++" strong binding; "nd" not determined. § peptide is not soluble in water.

Example IV: Identification of T cell epitopes by measuring cytokine responses to 15-mer peptide pools arrayed as matrix and confirmation of individual peptides

547 peptides, each individual peptide consisting of 15 aa and overlapping its precursor by 14 out of 15 aa, were designed to span the complete sequence of the CMV pp65 antigen. Sequences of all peptides are shown in Table 2.

Because the number of available human PBMC for screening T cell responses is usually limited, peptides were not applied individually, but combined to either column-pools or row-pools arrayed in a matrix format (Fig. 6). First, individual peptides were dissolved in 100% DMSO at 5 mg/ml. Then mixtures of 21 peptides each were prepared as such that each individual peptide was contained in exactly 2 pools. For restimulation of human PBMCs peptide mixtures were diluted in assay medium such that each peptide was present at a final concentration of 5µg/ml. This concentration is well above the concentration of 1,75 µg/ml, which Maecker et al. (Maecker paper) found to be saturating for CD4 and CD8 positive T cell responses. The final concentration

of DMSO in each sample was 2,1%.

Altogether PBMC from 10 healthy, CMV seropositive, HLA-A2 positive donors were used for screening of the 42 peptide pools represented in Fig. 6. To verify, whether class I epitopes usually shorter than 15 aa were properly recognized within the 15mer peptides and that the high DMSO content did not impair T cell activation and function, a single well-characterized immunodominant CD8-restricted 9-mer epitope from CMV pp65 (NLVPMVATV, pos. 495-503) was included in each screen at a concentration of 10 µg/ml (final DMSO concentration 0.1%).

As a read out for T cell reactivation a human IFN-gamma Elispot was used. Figure 8a shows a donor reacting with very few peptide pools. The crossover points of the positive column and row pools identify peptides 490-492. These 3 peptides contain the well-characterized immunodominant CD8-restricted 9-mer minimal epitope NLVPMVATV, pos. 495-503. Responses elicited by the peptide pools were largely equal to the response elicited with 9-mer minimal epitope (Fig. 8a). This indicates, that shorter epitopes are properly recognized within 15-mer peptides and that the high DMSO content did not impair the T cell response.

Most other donors showed a more complex response directed against several epitopes. One example is shown in figure 8b. The results of this primary screen with all 10 donors are summarized in Table 4.

Table 4. Summary of primary screen

Donor number	HLA type	Reacting with horizontal mixtures	Reacting with vertical mixtures	Defining the following peptides for retesting
9936	A2/24,B44/41	19	36,37,38,39	490,491,492,493
10511	A2/28,B16/40	2, 10, 19, 20, 21	31,33,34,35,37,38,40,41,42	37,39-41,43,44,46-48; 256,258-260,262,263,265-267; 485,487-489,491,492,494-496; 506,508-510, 512, 513, 515-517; 527,529-531,533,534;536-538
10632	A2/28,B12/27	19	29-32;34-39	483-486; 488-493
10687	A2/11,B7/13	10,11,17,18,19	22,23;33-42	247,248,258-267; 268,269,279-288; 394,395,405-414; 415,416,426,427,468-475;476,477,487-496
10689	A2/25,B13/18	8,19	22,23;28,29,30,36-41	205,206;211-214, 219-224;476,477;482-484;490-495
10736	A2/3,B15/35	19	35-38	489-492
10764	A2/24,B7/27	10,17,18,20	22;28-31;36-42	247,253-256,261-267;394,400-403,408-414;415,421-424;469-475;497,503-506;511-517

Donor number	HLA type	Reacting with horizontal mixtures	Reacting with vertical mixtures	Defining the following peptides for retesting
10788	A2/29,B44/60	2,3,4,14-16,17-19,21	22-42	28-32,36-48;49-53,57-105;331-335,339-351;352-356,360-372;373-377;381-393;394-398,402-414;415-419,423-427;468-475;476-480,484-496;518-522;526-538
10791	A2/3,B7/27	11,12,19	32-42	278-287;299-308;486-495

In a secondary screen all individual peptides corresponding to crossover points of positive row and column pools were retested. PBMC of the same donors were in this case incubated with the individual peptides at a final concentration of 20 µg/ml for 20hrs. Again a human IFN-gamma Elispot assay was used as a read out for T cell reactivation.

Fig. 9 shows the result from donor 10736, who had been identified in the primary screen to be an individual with a highly focused T cell response (Fig 8a). Retesting with individual peptides no. 489-495 confirmed reactivity of peptides 490-492 and in addition also identified peptide 493 as inducers of IFN-gamma secretion. The extent of the response to the 15mer peptides was again comparable to that of the 9-mer minimal epitope. A random selection of peptides corresponding to crossover points of negative row and column mixtures in the primary screen were included as negative controls; none did induce any IFN-gamma secretion (Fig. 9).

The complete results of this secondary screen and all epitopes defined by it are summarized in Table 5 (epitopes already described in the literature) and Table 6 (novel epitopes identified through the approach according to the present invention).

Among the already known epitopes, strikingly, the well-characterized CD8-restricted 9-mer minimal epitope (NLVPMVATV, pos. 495-503) was recognized in 9 out of 10 donors, proving that it is one of the most frequently recognized HLA-A*0201 restricted epitopes in the CMV pp65 antigen. Among the 10 HLA A2 donors 2 also expressed HLA-B7. From the literature 2 HLA-B7 restricted epitopes are known: pp65 265-274 and pp65 417-426. Both were found in both HLA B7 positive donors and especially the pp65 417-426 epitope seemed to induce even more IFN- γ secretion than the HLA-A*0201 restricted pp65 495-503 epitope. This data are in agreement with the observation, that T cell responses against the CMV pp65 antigen is strongly linked to expression of HLA-A2 and/or HLA-B7 epitopes (Eur. J. Immunol., 2000, 30, 2531-2539. Beside these so-called "immunodominant" epitopes most other described epitopes within pp65 were re-discovered (Table 5). This demonstrates, that the present approach is both very sensitive and comprehensive and thus suitable for detection of both immunodominant and subdominant epitopes.

Table 5a. Comparison of found epitopes with literature known class I epitopes

Peptides reactive in our screen	Peptide as described in literature	Described as	Described HLA restriction	literature	Frequency in screen ¹⁾
0	14-22	T cell epitope	A0201	Solache et al., 1999	0/9
# 10788: 117,118,119,120	120-128	T cell epitope	A0201	Solache et al., 1999	1/9
All except 10764: 489-493	495-503	T cell epitope	A0201	Wills et al., 1996 & Weekes et al., 1999a	8/9
#10764: 522	522-530 517-531	predicted T cell response	A0201	Solache et al., 1999 Kern et al., 1999	1/9
#10788: 108	110-118 109-123	Binding T cell response	A0201	Solache et al., 1999 Kern et al., 1999	1 / 9
#10687: 282,284 # 10791: 282, 283, 284?, 285, 287	286-295 289-303	Binding T cell response	A0201	Solache et al., 1999 Kern et al., 1999	2/9
#10511: 516?, 517 # 10764 513-517	519-527 517-531	Binding T cell response	A0201	Solache et al., 1999 Kern et al., 1999	2/9
0	316-338	T cell epitope	A0201	Bitmansour et al., 2001	0/9
Found: 6 out of 8 literature known epitopes in 9 HLA A2 positive donors					
0	353-375	T cell epitope	A1	Retiere et al., 2000	0/0
Found: 0 out of 1 literature known epitopes in 0 HLA A1 positive donors					
0	113-121	T cell epitope	A24	Retiere et al., 2001	0/2
0	328-337	T cell epitope	A24	Kuzushima et al., 2001	0/2
Found: 0 out of 2 literature known epitopes in 2 HLA A24 positive donors					
Possibly recognized in #10632	512-520	T cell epitope	B12	Wills et al., 1996& Weekes et al., 1999a	1/1

Found: 1 out of 1 literature known epitopes in 1 HLA B12 positive donor					
0	123-131	T cell epitope	B35	Gavin et al., 1993 & Wills et al., 1996	0/1
0	187-195	T cell epitope	B35	McLaughlin-Taylor et al., 1994	0/1
# 10687(B7 and not B35) : 394,395	397-411	T cell epitope	B35	Wills et al., 1996 & Weekes et al., 1999a	1/7
Found: 0 out of 3 literature known epitopes in 1 B35 positive donor; 1 found in HLA B35 negative donor					
#10687 : 260-265 #10764 : 261-264	265-274	T cell epitope	B7	Weekes et al., 1999b	2/2
#10687 : 414,415,416 #10764 : 413,414,415	417-426	T cell epitope	B7	Weekes et al., 1999a	2/2
Found: 2 out of 2 literature known epitopes in 2 HLA B7 positive donors					

1) Number of refinding/number of donors' expressing the respective HLA type

Table 5b. Comparison of found epitopes with literature known class II epitopes

Peptides re-active in our screen	Peptide as described in literature	Described as	Described HLA restriction	literature	Frequency in screen ¹⁾
0	34-56	T cell epitope	DQ0602	Bitmansour et al., 2001	0
0	364-386	T cell epitope	D1302	Bitmansour et al., 2001	0
#10788: 361	361-376	T cell epitope	DR11	Khattab et al., 1997	1
#10511 509,510	509-523	T cell epitope	DR3	Khattab et al., 1997 & Bitmansour et al., 2001	1
0	144-166	T cell epitope	DR1401	Bitmansour et al., 2001	0
0	177-191	T cell epitope	class II	Bitmansour et al., 2001	0
Possibly: #10687: 284 #10791: 285	285-299	T cell epitope	class II	Bitmansour et al., 2001	2
0	417-431	T cell epitope	class II	Bitmansour et al., 2001	0
Possibly: 10511: 487,489 10687: 487,488,489	489-503	T cell epitope	class II	Bitmansour et al., 2001	2
0	205-219	T cell epitope	?	Kern et al., 1999	0
Possibly #10687: 287,288	293-307	T cell epitope	?	Kern et al., 1999	1

1) no information on the MHC class II phenotype of the donors' is available

Table 6 lists all novel T cell epitopes which have been identi-

fied through the approach according to the present invention as described in Examples I and II.

The first group of epitopes (peptides 262, 394, 395, 411, 415, 416, 417) comprises sequences, which have already been described in the literature as T cell epitopes, albeit with a different HLA restriction than found here. The present results show that these epitopes are also recognized by T cells in a different HLA context than known before. This finding is new and expands the usefulness of these peptides e.g. as part of vaccines suitable for individuals expressing these particular HLAs.

The second group of epitopes (peptides 213, 214 and 216) have already been described in the literature as binding to HLA-A*0201 molecules, however before our work there was no data proofing that these peptides can activate T cells. Binding of a peptide is a pre-requisite for activation of/recognition by T cells but does not guarantee it. First, not all possible peptide sequences within an antigen are generated equally well or at all during antigen processing and presentation in vivo. Thus a synthetic peptide binding to HLA is not necessarily a naturally occurring epitope. Even if T cells can be primed against such peptides by active vaccination, they are not useful against the pathogen, because an infected cell does not display these peptides on its surface. Second, even if a binding peptide corresponds to a naturally occurring epitope this does not guarantee, that it can induce functional T cells. Instead certain peptides can act as antagonists and anergize T cells. The present data show for the first time that these peptides can induce functional, IFN-gamma secreting T cells in humans as a consequence of viral infection. This finding is new and provides a rationale for including these peptides in vaccines against CMV.

The third group of epitopes (peptides 211, 476, 477, 479, 421, 422, 423, 424, 469, 470, 503, 506) have so far not been described at all. Since the present data show that these peptides can induce functional, IFN-gamma secreting T cells in humans as a consequence of viral infection, they represent by themselves or contain within their sequence novel T cell epitopes. These are especially useful for inclusion in vaccines against CMV.

Since the screening described in this Example uses whole PBMCs, the complete T cell response to any of the 15mers was analyzed. This response can be restricted to any of the HLA-alleles expressed by the respective donors. These include, in each case at least two possible alleles of, HLA-A, -B, -C (class I) and HLA-DR, -DP, -DQ (class II). For further characterizing the novel epitopes provided herewith, one may define the exact HLA restriction of these epitopes and the minimal epitopes within the sequences recognized by T cells. Both can be done by a variety of well-established approaches known to the one skilled in the art (Current Protocols in Immunology, John Wiley & Sons, Inc.).

First, publicly available programs can be used to predict T cell epitopes on the basis of binding motifs. These include for instance: http://bimas.dcrt.nih.gov/molbio/hla_bind/ (Parker et al. 1994), <http://134.2.96.221/scripts/MHCServer.dll/home.htm> (Rammensee et al. 1999), <http://mypage.ihost.com/usinet.hamme76/> (Sturniolo et al. 1999). The latter prediction algorithm offers the possibility to identify promiscuous T helper-epitopes, i.e. peptides that bind to several HLA class II molecules. The respective HLA molecules, which were predicted with highest probability, are listed in Table 4. These prediction can be verified by testing of binding of the peptide to the respective HLA. Sequences, which were predicted to be possible class II epitopes, were tested for their binding to DRB1*0101, *0401 and *0404 molecules (see example III). The peptides 421, 469 and 470 (spanning aa 421-438, 469-484 and 470-485) were found to bind to DRB1*0404 (see Table 5)

A way of quickly discerning whether the response towards a peptide is class I or class II restricted is to repeat the ELISpot assay with pure CD4+ or CD8+ T cell effector populations. This can for instance be achieved by isolation of the respective subset by means of magnetic cell sorting. Pure CD8+ T cells can also be tested in ELISpot assays together with artificial antigen-presenting-cells, expressing only one HLA molecule of interest. One example are HLA-A*0201 positive T2 cells (174CEM.T2, Nijman et al., 1993). Alternatively, one can use ELISpot assays with whole PBMCs in the presence of monoclonal antibodies spe-

cifically blocking either the CD4+ or CD8+ T cell sub-population. Exact HLA restriction can be determined in a similar way, using blocking monoclonal antibodies specific for a certain allele. For example the response against an HLA-A24 restricted epitope can be specifically blocked by addition of an HLA-A24 specific monoclonal antibody.

For definition of the minimal epitopes within the peptide sequences recognized by T cells, one can e.g. synthesize series of overlapping and truncated peptides (e.g. 8-, 9-, 10-mers and retest these individually.

Example V: Confirmation of novel HLA-A*0201 epitopes by T2-ELISpot

In order to confirm predicted HLA-A2 epitopes within positive 15mers, 9- or 10-mer peptides were synthesized and tested individually. ELISpot assays were performed with T2 cells as antigen-presenting cells and isolated CD8+ T-cells from several donors. In this setting only optimal-length epitopes binding from the outside to HLA-A*0201 can trigger IFN- γ secretion. Five novel HLA-A*0201 epitopes could be confirmed by this approach (Table 6). The assay was carried out as described (Herr 1997). Briefly, T2 cells (Nijman 1993) were grown in ELISpot medium (see M&M section) and adjusted to 4×10^5 /ml 3 days prior use. After washing 2 times (PBS; 1% human albumin from SIGMA), cells were seeded at 2.5×10^6 /ml in ELISpot medium and 10 μ g/ml peptide was added for overnight culture. The next day T2 cells were washed once with ELISpot medium and 100 μ l (1×10^6 T2 cells/ml) were seeded into coated and blocked ELISpot plates (see above), supplemented with 20 μ g/ml peptide. For isolation of CD8+ lymphocytes 1×10^7 PBMC were mixed with 20 μ l anti human CD8+ Microbeads (Miltenyi Biotec, Bergisch Gladbach, Germany) in a final volume of 100 μ l and incubated for 15 min on ice. Afterwards cells were washed with a 10 times excess of MACS buffer and CD8+ cells were isolated on magnetic Minicolumns (Miltenyi). 100 μ l of CD8+ lymphocytes adjusted to 1×10^6 c/ml were added per well. After a culture period of 20 h cells were removed by washing 4 times with PBS, 0.05% Tween 20 and the assay was developed as described in the M&M section.

Table 6: CD8+ T-cell responses against HLA-A*0201 restricted epitopes of CMV pp65 assessed by T2 ELISpot assays for IFN- γ .

Donor ¹⁾	10511	10632	10687	10689	10788
Epitope ²⁾					
ILKEPVHGV (HIV, negative control)	17(\pm 4) ³⁾	16(\pm 3)	25(\pm 7)	77(\pm 2)	34(\pm 2)
NLVPMVATV ₄₉₅₋₅₀₃ (CMV, positive control)	<	165(\pm10)	193(\pm1)	189(\pm10)	172(\pm2)
RLLQTGIHV ₄₀₋₄₈	<	<	<	<	67(\pm9)
VIGDQYVKV ₂₁₈₋₂₂₆	<	31(\pm1)	112(\pm5)	<	100(\pm8)
YLESFCEDV ₂₂₇₋₂₃₅	<	<	<	<	<
AMAGASTSA ₄₂₅₋₄₃₃	<	<	96(\pm5)	<	104(\pm8)
FTWPPWQAGI ₄₈₂₋₄₉₁	<	<	<	<	70(\pm)
KYQEFFWDA ₅₀₉₋₅₁₇	<	32(\pm1)	<	<	72(\pm22)
RIFAELEGV ₅₂₂₋₅₃₀	<	<	92(\pm8)	<	71(\pm18)

¹⁾ new blood draws 3-6 months apart from primary matrix screen

²⁾ new epitopes shown boldface

³⁾ spot numbers per 100,000 CD8+ T-cells; <: below background (spots with HIV negative control peptide) plus 2 standard deviations.

Example VI: Confirmation of novel class II (HLA-DR4) epitopes identified by the Epitope Capture Method using human PBMC

Among others, 15mers 55-61 were identified by the epitope capture method as strong binders of soluble HLA-DRB1*0401 (Fig. 7a, 7b). Binding of individual overlapping 15mers 55-61 identifies the amino acid sequence YTPDSTPCH as core binding region of LILVSQYTPDSTPCHRGDNQL which may contain several versions of a class II epitope. It is well known that class II epitopes have ends of variable length protruding the MHC binding groove. Interestingly, donor 10788 showed strong T-cell responses against 15mers 57 and 59 (Fig. 10) measured by IFN- γ ELISpot as described

in Example IV. This confirms that LILVSQYTPDSTPCHRGDNQL represents or contains at least one HTL epitope binding to at least HLA-DRB1*0401 originally discovered by the epitope capture method of the present invention.

Fig. 10: PBMNC from subject 10788 were applied for IFN- γ ELISpot with CMVpp65 15mers 57, 59 and controls (med: no peptide, HIV: irrelevant HIV-derived peptide, ConA: polyclonal stimulation)

As a further example 15mers 469 and 470 identified by the epitope capture method of the present invention as weakly binding to DRB1*0404 (Tab. 3) were tested for reactivity with human PBMNC. To distinguish between CD8-positive CTL mediated or CD4-positive HTL mediated reactivity, intracellular cytokine staining combined with staining for surface markers was performed: after thawing, PBMNC were kept overnight (37°C, 5% CO₂) in RPMI1640 with 10% human serum type AB (BioWhittaker). Next day aliquots of 2 million PBMNC were incubated with peptide (80 μ g/ml) or Concanavalin A (SIGMA). After 1h, 10 μ g/ml brefeldin A (SIGMA) were added and incubation was continued for 5 hours. Surface staining for CD4 (phycoerythrin), CD8 (cyochrome), CD69 (allophycocyanine) was done with antibodies labelled as indicated (Pharmingen, San Diego, CA, USA). After fixation with 1% para-formaldehyde in PBS, cells were permeabilized by incubating for 15 min in 0.5% BSA/0.1% Na-azide/0.1% saponin in PBS. Intracellular cytokines were stained with anti-IFN- γ (FITC) antibody 4S.B3 (Pharmingen). Samples (100,000 events in the lymphocyte gate) were read on a FACScalibur (Becton Dickinson).

15mers 469 and 470 covering DEDSDNEIHNPVFTW₄₆₉₋₄₈₄ contain both A*0201 and class II binding motifs and bind to soluble DRB1*0404 (Tab. 3). Intracellular cytokine staining confirmed significant IFN- γ secretion in both the CD8 and CD4 positive T-cell compartments for donor 10788 (Fig 11, lower two panels). Conversely, donor 10687 showed only CD8 mediated IFN- γ secretion against 15mer 489 AGILARNLVPMVATV₄₈₉₋₅₀₃ indicating that in this case only the HLA-A2 epitope NLVPMVATV₄₉₅₋₅₀₃ was targeted. These results confirm that peptide DEDSDNEIHNPVFTW₄₆₉₋₄₈₄ contains both class I and class II restricted epitopes.

Fig. 11: Confirmation of simultaneous CD4+ and CD8+ T-cell responses against CMVpp65 15mers 469, 470 by intracellular IFN- γ staining. PBMC from 2 donors (10687, 10788) were stimulated with either ConA as positive control (1st column), 15mers containing both putative class I and class II epitopes (columns 2 and 3) or medium as negative control (right column). Cells were stained for intracellular IFN- γ (x-axis) or surface the T-cell differentiation markers CD4 or CD8 (y-axis). Percentage in upper-right quadrant is indicated, numbers significant over background are shown boldface.

Example VII: Confirmation of novel class II (HLA-DRB1*0401) epitopes identified by the Epitope Capture Method using HLA-DR4 transgenic mice

For those, DRB1*0401 binding, peptides where T cell reactivity was not demonstrated by testing CMV seropositive individuals, experiments in HLA-DR4 transgenic mice, expressing DRB1*0401 molecules, were performed. The longer peptides (no. 1500-1505), covering all candidate epitopes binding to DRB1*0401 molecules, were synthesized (see Table 7) and injected into the mice. One week after the last injection total murine splenocytes were re-stimulated ex vivo with the same peptides that were used for vaccination, as well as with overlapping 15-mers, representing corresponding longer peptides and an irrelevant, influenza hemagglutinin derived peptide (no. 1171) as a negative control. T cell reactivity was determined by INF-gamma ELISpot assay (Fig. 12) and as a result, five out of six longer peptides were shown to be immunogenic. Moreover, most of the 15-mers, representing longer peptides and showing affinity to DRB1*0401 molecules, were also verified to re-activate ex vivo T cells from DRB1*0401-transgenic mice. One peptide (no.1503) did not induce immune response at least under the immunization conditions used in these studies.

For the fine epitope mapping and in vivo testing of CD4+ specificity of immunogenic peptides, mice were vaccinated with the longer peptide and splenocytes were separated into CD4+ and CD8+ T cell populations (92-94 % purity for CD4+ fraction) to be tested for IFN- γ production as described above. Fig. 12 shows the

results of such ELISpot assay. In all cases the T cell response measured with splenocytes could be confirmed using CD4+ cells. Usually the CD8+ response was negligible. A simultaneous CD4+ and CD8+ response as seen for peptide 1502 could be due to an additional class I mouse epitope contained within the peptide sequence.

Table 7a summarizes all the data of peptide binding studies with soluble DR molecules and mouse experiments. The good correlation between peptide affinities to DRB1*0401 molecules and specific stimulation of CD4+ cells in HLA-DR4 transgenic mice was observed. For instance, peptides no.1500-1502 and 1504 showed similar results in both approaches. Peptide no. 1505 that was not so good in binding assay evoked the highest frequency of T cells generated in mice after peptide injection. Peptide no. 1503 that had weak affinity to DRB1*0401 molecules in vitro was not immunogenic in mice injected either with CpG1668 or with CFA/IFA. Here should be also mentioned that some of peptides showed variable immunogenicity and specific CD4+ frequency depending on the adjuvant used for the injection. Based on the results of mouse ELISpot assay with enriched fraction of CD4+ cells and peptide binding assay with soluble DRB1*0401 molecules, epitope core sequences recognized by DRB1*0401 carrying T cell (see Table 7 and Table 7a) can be determined. Four of them fit well to the predicted by TEPITOPE algorithm sequences, except one for peptide no.1501 which is not predicted to bind DRB1*0401 molecules.

Table 7. Verification of DRB1*0401 ligands, as candidate epitopes, in transgenic mice.

Proposed DRB1*0401 core binding motifs are highlighted.

No.	Peptide sequence	AA	Binding to DRB1*					DRB1*0401 mice		Comment, reference
			0401	0404	0101	0701	1101	total SC	CD4+	
1500	PSLILVSQYTPDSTPCHR-GDNQLQVQHTR	53-81	++	++	+	-	-	+	+	new epitope, human PBMC's in this study
1501	QWKEPDVYVTSFVFPTK-DVALR	174-196	+	-	++	-	+/-	+	+	Bitmansour, et al., CD4+ clone
1502	LLQRGPOYSEHPT-FTSQYRIQG	356-377	+	-	-	-	++	+	++	Khattab et al., CD4+ clone/DR11
1503	YRHTWDRHDEGAAGG-DDDVW	381-400	+/-	-	-	-	-	-	-	not immunogenic
1504	TSGVMTRGRKKAES-FVAPEEDTDE	447-470	+	+++	+	-	+	+	++	new epitope
1505	QGQN-LKYQEFFWDANDIYRIF	504-524	+/-	-	-	-	-	++	++	Khattab et al., CD4+ clone/ DR3

"-" no binding/T cell response; "+/-" weak binding/T cell response; "+" intermediate binding/T cell response; "++" strong binding/T cell response

In conclusion, most of the peptides discovered as ligands for soluble DRB1*0401 molecules in the present "reverse immunological" approach were confirmed to elicit specific CD4+ response in DRB1*0401-transgenic mice. Thus, five out of six epitope candidate are real DRB1*0401 epitopes. Additionally, they are promiscuous (see Table 7 and Table 7a). Three of them (no. 1501, 1502 and 1503) have been described in the literature earlier but not as DRB1*0401 specific epitopes. Two DRB1*0401 binders (no. 57 and 59) and three peptides binding to DRB1*0404 (no. 421, 469 and 470) were also found to induce INF-gamma production in PBMC's from CMV seropositive donors (see Example IV and VI). All together, it indicates that the present Epitope Capture Approach can identify true T cell epitopes.

Table 7a. Summary of identification of human class II epitopes using soluble DRB1*0401 molecule molecule and HLA DR4-transgenic mice.

lighted.

Pep no.	Peptide sequence	Binding to DRB1					DRB1*0401 mice		Comments
		*0401	*0404	*0101	*0701	*1101	total SC	CD4+	
1500	PSLILV <u>SQYTPD</u> STPCHRG-DNQLQVQHTR	++	++	+	-	-	+	+	AA 53-81, newly identified
54	SLILV <u>SQYTPD</u> STPC	-					-	+/-	
55	LILV <u>SQYTPD</u> STPCH	+					++	+	
56	ILV <u>SQYTPD</u> STPCHR	++					++	+	
57	L <u>V</u> SQYTPDSTPCHRG	++	-	-			++	+	
58	V <u>SQYTPD</u> STPCHRGD	++	-	+	-	-	++	+	
59	SQYTPDSTPCHRGDN	++					++	+/-	
60	QYTPDSTPCHRGDNQ	++					++	+/-	
61	YTPDSTPCHRGDNQL	+					+	+/-	
62	TPDSTPCHRGDNQLQ	-					-	-	
63	PDSTPCHRGDNQLQV	-					-	-	
64	DSTPCHRGDNQLQVQ	+					-	-	
65	STPCHRGDNQLQVQH	-					-	-	
1501	QWKEPDVYYTSAFVFETK-DVALR	+	-	++	-	+	+	+	AA 174-196, known class II
174	QWKEPDVYYTSAFVF	-					+/-	+/-	
175	WKEPDVYYTSAFVFP	-					+	+	
176	KEPDVYYTSAFVFPT	+/-					+	+	
177	EPDVYYTSAFVFPTK	+	-	++	-	+	+	+	
178	PDVYYTSAFVFPTKD	++					+	+	
179	DVYYTSAFVFPTKDV	+					+	+	
180	VYYTSAFVFPTKDVA	+/-					+	+/-	
181	YYTSAFVFPTKDVAL	-					+	+/-	

182	YTSAFVFPTKDVALR						+/-	+/-	
150 2	LLQRGPQYSEHPT- FTSQYRIQG	+	-	-	-	++	++	++	AA 356-377, known DR 11
356	LLQRGPQYSEHPTFT						+/-	+/-	
357	LQRGPQYSEHPTFTS	-					+	+/-	
358	QRGPQYSEHPTFTSQ	+					++	+/-	
359	RGPQYSEHPTFTTSQY	+/-	-	-	-	+/-	++	++	
360	GPQYSEHPTFTTSQYR	+/-					++	++	
361	PQYSEHPTFTTSQYRI	+/-					+	++	
362	QYSEHPTFTTSQYRIQ	-					+	++	
363	YSEHPTFTTSQYRIQG						+	++	
150 3	YRHTWDRHDEGAAGDDDW	+/-	-	-	-	-	-	-	AA 381- 400, not im- munogenic
380	EYRHTWDRHDEGAAG	-					-	-	
381	YRHTWDRHDEGAAGG	-					-	-	
382	RHTWDRHDEGAAGGD	-					-	-	
383	HTWDRHDEGAAGGDD	+/-					-	-	
384	TWDRHDEGAAGGDD	+/-	-	-	-	-	-	-	
385	WDRHDEGAAGGDDV	-					-	-	
150 4	TSGVMTRGRKKAES- TVAPEEDTDE	+	++	+	-	+	+	++	AA 447-470, newly iden- tified
448	RGVMTRGRKKAESTV	-					+/-	-	
449	GVMTRGRKKAESTVA	+/-					+	++	
450	VMTRGRKKAESTVAP	+					+	++	
451	MTRGRKKAESTVAPE	+					+	++	
452	TRGRKKAESTVAPEE	+	++	++	-	-	+	++	
453	RGRKKAESTVAPBED	+					+/-	+/-	
454	GRKKAESTVAPBEDT	+					+/-	+/-	
455	RLKAESTVAPBEDTD	-					-	-	
456	LKAESTVAPBEDTDE						-	-	

150 5	QGQNLKYQEFFWDANDIYRIF	+	-	-	-	-	++	++	AA 504-524, known DR3
504	QGQNLKYQEFFWDAN	-					-	-	
505	GQNLKYQEFFWDAND	-					+	-	
506	QNLKYQEFFWDANDI	-					++	+/-	
507	NLKYQEFFWDANDIY	+/-	-	-	-	-	++	+	
508	LKYQEFFWDANDIYR	+/-					++	++	
509	KYQEFFWDANDIYRI	-					++	++	
510	YQEFFWDANDIYRIF	-					++	++	

"-" no binding/T cell response; "+/-" weak binding/T cell response; "+" intermediate binding/T cell response; "++" strong binding/T cell response

References

1. Aichinger G et al., 1997. J. Biol. Chem. 272: 29127-29136.
2. Ausubel FM et al. (editors), 2001. Current Protocols in Molecular Biology, Volumes 1-4, John Wiley and Sons (publishers).
3. Bitmansour AD et al., 2001. J. Immunol. 167: 1151-1163.
4. Bonifacino JS et al. (editors), 2001. Current Protocols in Cell Biology, Volumes 1-2, John Wiley and Sons (publishers).
5. Britt WJ et al, 1996. Chapter 77 "Cytomegalovirus", Virology by Fields et al. (editors), Livincott-Raven Publishers
6. Cox AL et al., 1994. Science 264: 716-719
7. Coligan JE et al. (editors), 2001. Current Protocols in Immunology, Volumes 1-4, John Wiley and sons (publishers).
8. Drew WL et al, 1999. Curr Clin Top Infect Dis 19: 16-29
9. Field AK et al., 1999. Antivir Chem Chemother 10(5): 219-232
10. Fowler KB et al, 1992. N Engl J Med 326: 663-673
11. Gallot G. et al., 2001. J. Immunol. 167: 4196-4206.
12. Gavin MA et al., 1993. J Immunol. 151: 3971-80
13. Gorga, JC et al., 1987. J. Biol. Chem. 262: 16087-16094.
14. Greenberg, P. D. et al, 1999. Science 285: 546.
15. Heemels, MT et al, 1995. Annu. Rev. Biochem. 64: 463-491
16. Kern F et al, 2000. Eur J Immun 30: 1676-1682
17. Kern F et al, 1999. J Virol 73(10): 8179-8184.
18. Khat tab BA et al., 1997. J Med Virol 52: 68-76.
19. Klein, J, 1986. Natural History of the MHC, John Wiley and Sons (publishers)
20. Komanduri, KV et al., 1998. Nat. Med. 4: 953.
21. Kronenberg M et al., 1999. Immunol Today 20: 515-21.
22. Kuzushima K et al., 2001. Blood 98(6): 1872-1880.
23. Kwok WW et al., 2001. Trends Immunol 22(11): 583-588.
24. Lalvani A et al., 1997. J. Exp. Med. 186 (6): 859-865.
25. McLaughlin-Taylor E et al., 1994. J Med Virol 43: 103-110.
26. Maecker HT et al, 2001. J Immunol Methods 255: 27-40
27. Masuoka M et al., 2001. Viral Immunology 14(4): 369-377
28. Morgan CL et al., 1998, Immunogenetics 48(2): 98-107.
29. Nichols WG et al, 2000, J Clin Virol 16(1): 25-40
30. Nijman HW et al, 1993. Eur. J. Immunol., 6, 1215-9
31. Novak N et al., 2001. J. Immunol. 166: 6665-6670.
32. Parker, KC et al., 1994. J. Immunol. 152: 163.
33. Plotkin SA, 1999. Pediatr Infect Dis J 18(4): 313-25

34. Rammensee, HG et al., 1999. Immunogenetics 50: 213-219
35. Reddehase MJ, 2000. Curr Opin Immunol. 12: 390-396
36. Retriere C et al, 2000. J Virol 74(9): 3948-3952
37. Saulquin X et al., 2000. Eur J Immunol 30 : 2531-2539
38. Sia IG et al, 2000. Clin Microbiol Rev 13(1) : 83-121
39. Solache A et al., 1999. J. Immunol. 163: 5512-5518.
40. Stern LJ et al., 1994. Structure 2: 245-251
41. Sturniolo T et al., 1999. Nature Biotechnology 17: 555-562.
42. Tana et al., 1998, J. Human Genet. 43(1): 14-21.
43. Tobery WT et al., 2001. J Immunol Methods 254: 59-66.
44. Valli A et al., 1993. J. Clin. Invest. 91: 616-628
45. Van den Eynde BJ, et al., 1997. Curr Opin Immunol. 5: 684-93
46. van Son WJ, 1999. Intervirology 42(5-6): 285-290
47. Villadangos JA et al., 2000. Immunity 12: 233-239
48. Waldrop SL et al., 1998. J. Immunol. 161: 5284.
49. Wilson DB et al., 1999. J. Immunol. 163: 6424-6434.
50. Weekes MP et al., 1999a. J Immunol 162: 7569-7577.
51. Weekes MP et al., 1999b, J Virol 73(3): 8140-8150.
52. Zaia JA et al, 2000. Cytomegalovirus Prevention and Treatment in 2000. Hematology (Am Soc Hematol Educ Program) 2000: 339-355

Claims:

1. Method for isolating ligands which have a binding capacity to a MHC/HLA molecule or a complex comprising said ligand and said MHC/HLA molecule characterised by the following steps:
 - providing a pool of ligands, said pool containing ligands which bind to said MHC/HLA molecule and ligands which do not bind to said MHC/HLA molecule,
 - contacting said MHC/HLA molecule with said pool of ligands whereby a ligand which has a binding capacity to said MHC/HLA molecule binds to said MHC/HLA molecule and a complex comprising said ligand and said MHC/HLA molecule is formed,
 - detecting and optionally separating said complex from the ligands which do not bind to said MHC/HLA molecule and
 - optionally isolating and characterising the ligand from said complex.
2. Method for isolating T cell epitopes which have a binding capacity to a MHC/HLA molecule or a complex comprising said epitope and said MHC/HLA molecule characterised by the following steps:
 - providing a pool of ligands, said pool containing ligands which bind to a MHC/HLA molecule and ligands which do not bind to said MHC/HLA molecule,
 - contacting said MHC/HLA molecule with said pool of ligands whereby a ligand which has a binding capacity to said MHC/HLA molecule binds to said MHC/HLA molecule and a complex comprising said ligand and said MHC/HLA molecule is formed,
 - detecting and optionally separating said complex from the ligands which do not bind to said MHC/HLA molecule,
 - optionally isolating and characterising the ligand from said complex,
 - assaying said optionally isolated ligand or said complex in a T cell assay for T cell activation capacity and
 - providing the optionally isolated ligand with a T cell activation capacity as T cell epitope or as complex.
3. Method according to claim 1 or 2, characterised in that said pool of ligands is selected from the group consisting of a pool of peptides, especially overlapping peptides, a pool of protein

fragments, a pool of glycolipids, a pool of glycosphingolipids, a pool of lipopeptides, a pool of lipids, a pool of glycans, a pool of modified peptides, a pool obtained from antigen-presenting cells, preferably in the form of total lysates or fractions thereof, especially fractions eluted from the surface or the MHC/HLA molecules of these cells, a pool comprised of fragments of cells, especially pathogen cells, tumor cells or tissues, a pool comprised of peptide libraries, pools of (poly)-peptides generated from recombinant DNA libraries, especially derived from pathogens or tumor cells, a pool of proteins and/or protein fragments from a specific pathogen or mixtures thereof.

4. Method according to any one of claims 1 to 3, characterised in that said MHC/HLA molecules are selected from HLA class I molecules, HLA class II molecules, non classical MHC/HLA and MHC/HLA-like molecules or mixtures thereof, or mixtures thereof.

5. Method according to any one of claims 1 to 4, characterised in that said characterising of the ligands of the complex is performed by using a method selected from the group consisting of mass spectroscopy, polypeptide sequencing, binding assays, especially SDS-stability assays, identification of ligands by determination of their retention factors by chromatography, especially HPLC, or other spectroscopic techniques, especially violet (UV), infra-red (IR), nuclear magnetic resonance (NMR), circular dichroism (CD) or electron spin resonance (ESR), or combinations thereof.

6. Method according to any one of claims 1 to 5, characterised in that it is combined with a cytokine secretion assay, preferably with an Elispot assay, an intracellular cytokine staining, FACS or an ELISA.

7. Method according to any one of claims 1 to 6, characterised in that said T cell assay comprises the mixing and incubation of said complex with isolated T cells and subsequent measuring cytokine secretion or proliferation of said isolated T cells.

8. Method according to any one of claims 1 to 6, characterised in that said T cell assay comprises measuring up-regulation of

activation markers, especially CD69, CD38, or down-regulation of surface markers, especially CD3, CD8 or TCR.

9. Method according to any one of claims 1 to 8, characterised in that said T cell assay comprises measuring up-/down-regulation of mRNAs involved in T cell activation, especially by real-time RT-PCR.
10. Method according to any one of claims 1 to 8, characterised in that said T cell assay is selected from T cell assays measuring phosphorylation/de-phosphorylation of components downstream of the T cell receptor, especially p56 lck, ITAMS of the TCR and the zeta chain, ZAP70, LAT, SLP-76, fyn, and lyn, T cell assays measuring intracellular Ca⁺⁺ concentration or activation of Ca⁺⁺-dependent proteins, T cell assays measuring formation of immunological synapses, T cell assays measuring release of effector molecules, especially perforin, granzymes or granulolysin or combinations of such T cell assays.
11. T cell epitopes identifiable by a method according to any one of claims 2 to 10, said T cell epitopes being selected from the group consisting of polypeptides comprising the sequence KM-QVIGDQYV, FTWPPWQAGI, AMAGASTSA, SDNEIHNPV, KYQEFFWDA or combinations thereof.
12. HLA A0201 binding epitopes with T cell activating capacity identifiable by a method according to any one of claims 2 to 10 using HLA A0201 molecules as MHC/HLA molecules, said HLA A0201 binding epitopes being selected from the group consisting of polypeptides comprising the sequence RLLQTGIHV, VIGDQYVKV, YLES-FCEDV or combinations thereof.
13. Use of a peptide comprising the sequence RPHERNGFTV for preparing a composition for activating T cells in an individual being B7-negative.
14. Use of a peptide comprising the sequence DDVWTSGSDSDE for preparing a composition for activating T cells in an individual being B35-negative.

15. Use of a peptide comprising the sequence TPRVTGGGAM for preparing a composition activating T cells in an individual being B7-negative.
16. Peptide binding to class II HLA molecules selected from peptide nos. 55-64, 109, 383, 384, 421, 449-454, 469 and 470 according to table 3.
17. Epitope or peptide according to any one of claims 11 to 16 characterized in that it further comprises 1 to 30, preferably 2 to 10, especially 2 to 6, naturally occurring amino acid residues, especially at the N-terminus, the C-terminus or at the N- and C-terminus.
18. Epitope or peptide according to any one of claims 11 to 17, characterised in that it further comprises a non-naturally occurring amino acid(s), preferably 1 to 1000, more preferred 2 to 100, especially 2 to 20 non-naturally occurring amino acid residues, at the N-terminus, the C-terminus or at the N- and C-terminus.
19. Use of an epitope or peptide according to any one of claims 11 to 18 for the preparation of a HLA restricted vaccine for treating or preventing cytomegalovirus (CMV) infections.
20. Use of an epitope according to any one of claims 11, 17 or 18 for the preparation of a vaccine for treating or preventing cytomegalovirus (CMV) infections.
21. Vaccine for treating or preventing cytomegalovirus (CMV) infections comprising an epitope according to any one of claims 11, 17 or 18.
22. HLA specific vaccine for treating or preventing cytomegalovirus (CMV) infections comprising an epitope or peptide according to any one of claims 11 to 18.
23. Vaccine as defined in any one of claims 19 to 22, characterised in that it further comprises an immunomodulating substance, preferably selected from the group consisting of polycationic

substances, especially polycationic polypeptides, immunomodulating nucleic acids, especially deoxyinosine and/or deoxyuracile containing oligodeoxynucleotides, or mixtures thereof.

24. Vaccine as defined in any one of claims 19 to 23, characterised in that it further comprises a pharmaceutically acceptable carrier.

25. Vaccine as defined in any one of claims 19 to 24, characterized in that said epitope is provided in a form selected from peptides, peptide analogues, proteins, naked DNA, RNA, viral vectors, virus-like particles, recombinant/chimeric viruses, recombinant bacteria or dendritic cells pulsed with protein/peptide/RNA or transfected with DNA comprising the epitopes.

26. T cells, a T cell clone or a T cell population or preparation specifically recognizing an epitope or peptide according to any one of claims 11-18.

27. Use of T cells, a T cell clone or a T cell population or preparation according to claim 26 for identification of heteroclitic epitopes.

28. Use of T cells, a T cell clone or a T cell population or preparation according to claim 26 for the preparation of a composition for therapy of CMV patients.


29. Method according to any one of claims 1 to 10, characterised in that the ligands or fragments of antigens, especially antigens from human immune deficiency virus (HIV), hepatitis A and B viruses, hepatitis C virus (HCV), Rous sarcoma virus (RSV), Epstein Barr virus (EBV), Influenza virus, Rotavirus, Staphylococcus aureus, Chlamydia pneumoniae, Chlamydia trachomatis, Mycobacterium tuberculosis, Streptococcus pneumoniae, Bacillus anthracis, Vibrio cholerae, Plasmodium sp. (Pl. falciparum, Pl. vivax, etc.), Aspergillus sp. or Candida albicans, tumor antigens or autoimmune antigens.

30. Method according to any one of claims 1 to 10 and 29, char-

acterised in that said ligands are HCV-, HIV-, HAV-, HBV-, RSV-, EBV-, Influenza virus- or Rotavirus-peptides having a binding capacity for an MHC/HLA-molecule.


1/19

Fig. 1. Peptide binding affinities to soluble DR4 molecules




2/19

Fig. 2. Identification of peptides capable bind to „empty“ DR4 molecules.
A. Purification of HLA-peptide complexes. B. MS analysis of bound peptides.



3/19

Fig. 3. Binding of high affinity peptide in the presence of the excess of low affinity peptide to DR4 molecules



4/19

Fig. 4. Binding of the individual peptides and peptide mixtures to DR4 molecules



5/19




Fig. 5a

6/19

Elispot with isolated CD8+ T cells and HLA A0201 molecules
(costimulation with anti CD28 mab)




Fig. 5b

7/19

Column pools →
 Row pools →

	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42
1	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27
2	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48
3	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69
4	106	107	108	109	110	111	112	113	114	115	116	117	118	119	120	121	122	123	124	125	126
5	127	128	129	130	131	132	133	134	135	136	137	138	139	140	141	142	143	144	145	146	147
6	150	151	152	153	154	155	156	157	158	159	160	161	162	163	164	165	166	167	168	169	170
7	184	185	186	187	188	189	190	191	192	193	194	195	196	197	198	199	200	201	202	203	204
8	205	206	207	208	209	210	211	212	213	214	215	216	217	218	219	220	221	222	223	224	225
9	226	227	228	229	230	231	232	233	234	235	236	237	238	239	240	241	242	243	244	245	246
10	247	248	249	250	251	252	253	254	255	256	257	258	259	260	261	262	263	264	265	266	267
11	268	269	270	271	272	273	274	275	276	277	278	279	280	281	282	283	284	285	286	287	288
12	289	290	291	292	293	294	295	296	297	298	299	300	301	302	303	304	305	306	307	308	309
13	310	311	312	313	314	315	316	317	318	319	320	321	322	323	324	325	326	327	328	329	330
14	331	332	333	334	335	336	337	338	339	340	341	342	343	344	345	346	347	348	349	350	351
15	352	353	354	355	356	357	358	359	360	361	362	363	364	365	366	367	368	369	370	371	372
16	373	374	375	376	377	378	379	380	381	382	383	384	385	386	387	388	389	390	391	392	393
17	394	395	396	397	398	399	400	401	402	403	404	405	406	407	408	409	410	411	412	413	414
18	415	416	417	418	419	420	421	422	423	424	425	426	427	428	429	430	431	432	433	434	435
19	436	437	438	439	440	441	442	443	444	445	446	447	448	449	450	451	452	453	454	455	456
20	457	458	459	460	461	462	463	464	465	466	467	468	469	470	471	472	473	474	475	476	477
21	518	519	520	521	522	523	524	525	526	527	528	529	530	531	532	533	534	535	536	537	538

Fig. 6. CMV pp65 peptide pool array

8/19

Fig. 7a. Binding of peptide pools derived from pp65 protein to DR4 molecules




Fig. 7b. Binding of single pp65 peptides to DR4 molecules





Fig 8a


1st screen with with peptide mixtures
(containing 21 peptides each)



Donor #10736
HLA A 2/3; HLA B 15/35

Fig 8b

1st screen with with peptide mixtures
(containing 21 peptides each)




Donor #10687
(HLA A2/11; HLA B7/13)

12/19


Fig 9

2nd screen with with single peptides

Donor #10736
HLA A 2/3; HLA B 15/35



13/19

Fig. 10

14/19

Fig. 11









Fig. 12 A:


16/19

Fig. 12 B:

17/19

Fig. 12 C:

18/19

Fig. 12 D:

19/19

Fig. 12 E: